

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

# (12) UK Patent Application (19) GB (11) 2 300 859 (13) A

(43) Date of A Publication 20.11.1996

(21) Application No 9615814.2

(22) Date of Filing 17.01.1994

Date Lodged 26.07.1996

(30) Priority Data

(31) 9600686

(32) 15.01.1993

(33) GB

(62) Derived from Application No. 9513418.5  
under Section 15(4) of the Patents Act 1977

(71) Applicant(s)

Imperial Cancer Research Technology Limited

(Incorporated in the United Kingdom)

Sardinia House, Sardinia Street, LONDON,  
WC2A 3NL, United Kingdom

(72) Inventor(s)

Agamemnon Antoniou Epenetos

Robert Anthony Spooner

Mahendra Deonarain

(51) INT CL<sup>6</sup>

A61K 47/48, C07K 16/00 19/00, C12N 15/62, G01N  
33/80, // A61K 51/10, C12N 9/22 (C07K 19/00 C07M  
5:00)

(52) UK CL (Edition O)

C3H H87M H87P H650 H651 H654 H656 H657 H660  
H675 H684 H685 H686 H690 H730  
U1S S1313 S1332 S1333 S1334 S1337

(56) Documents Cited

WO 91/16069 A1

Cell Biophysics 1992,21,121-138

(58) Field of Search

UK CL (Edition O) C3H H87E H87M H87P

INT CL<sup>6</sup> A61K 47/48

ONLINE: WPI

(74) Agent and/or Address for Service

Eric Potter Clarkson

St Mary's Court, St Mary's Gate, NOTTINGHAM,  
NG1 1LE, United Kingdom

(54) Compounds comprising a target cell-specific portion (TCP) and a cytotoxic portion (CP)

(57) Compounds are described which are selected from:

(i) TCP, preferably an ScFv and/or one which binds selectively to a tumour cell, fused to an oligomeric ribonuclease, preferably mammalian seminal ribonuclease;


(ii) a oligomeric complex of at least two molecules, each comprising a TCP fused to a further (cytotoxic) portion, said molecules being complexed to one another via said further portion;

(iii) TCP, other than an antibody, comprising at least two binding sites for the target cell, linked to a further (cytotoxic) portion;

(iv) a recognition portion, preferably an ScFv, for 4-hydroxy-3-iodophenylacetic acid or 4-hydroxy-3-nitrophenylacetic acid bonded to CP, preferably the biotin-binding portion of streptavidin or one which has nucleolytic activity, especially the catalytic activity of either DNA endonuclease or RNAase, which may be bonded to TCP.

In compounds (ii) and (iii), the further (cytotoxic) portion may be mammalian seminal ribonuclease or a biotin-binding portion of streptavidin and the TCP preferably recognises either a blood group antigen (for use in the determination of a blood group by haemagglutination) or an ScFv antibody fragment. Nucleic acid sequences encoding compounds (i) to (iii) are disclosed.

The compounds are of use in therapeutic systems for the treatment of cancer.






Fig. 1(a)

Fig. 1(b)






Fig. 1(c)

Fig. 1(d)






FIGURE 2

3/38

VHBACK2                    L Q E S G P  
CAGGTGCAGCTGCAGGAGTCAGGACC    VHBACK2  
                         PstI

BAMLINKERFOR            G S  
GGATCCGACATCGAGCTCACTCAGTCTCCA  
                         BamHI

BAMVλBACK                G S Q A E L T Q E S  
AAGCTTGGATCCCAGGCTGTTGTGACTCAGGAATCT  
                         BamHI

ECOVλFOR                K L T V L G R S \* \*  
CCAAACTGACTGTCCTAGGTCTCGAGTAATAAGAATTCATGC  
                         XhoI                    EcoRI

VHBACK3                    L Q Q P G  
CAGGTCCAACCTGCAGCAGCCTGG  
                         PstI

VH1FOR-2                G Q G T T L T  
GGGGCCAAGGGACCACGGTCACCGTCTCCTCA  
                         BstEII

FIGURE 3

4/38  
30

10                      20                      40                      50

M K Y L L P

AAGCTTGCATGCAAATTCTATTTCAAGGAGACAGTCATATGAAATACCTATTGCCT  
HindIII SphI                      SD                      /-----/

60                      70                      80                      90                      100                      110

T A A A G L L L L A A Q P A M A Q V Q  
ACGGCAGCCGCTGGATTGTTACTCGCTGCCCAACCAGCGATGGCCAGGTGCAG  
-----pelB leader-----/VHNP

120                      130                      140                      150                      160                      170

L Q Q P G A E L V K P G A S V K L S C  
CTGCAGCAGCCTGGGGCTGAGCTTGTGAAGCCTGGGGCTTCAGTGAAGCTGTCCTGC  
PstI

180                      190                      200                      210                      220                      2

K A S G Y T F T S Y W M H W V K Q R P  
AAGGCTTCTGGCTACACCTTCACCAGCTACTGGATGCACTGGGTGAAGCAGAGGCCT  
CDR1

30                      240                      250                      260                      270                      280

G R G L E W I G R I D P N S G G T R Y  
GGACGAGGCCTTGAGTGGATTGGAAGGATTGATCCTAATAGTGGTGGTACTAAGTAC  
CDR2

290                      300                      310                      320                      330                      340

N E K F L S K A T L T V D K P S S T A  
AATGAGAAGTTCAAGAGCAAGGCCACACTGACTGTAGACAAACCCTCCAGCACAGCC

350                      360                      370                      380                      390                      40

Y M Q L S S L T S E D S A V Y Y C A R  
TACATGCAGCTCAGCAGCCTGACATCTGAGGACTCTGCGGTCTATTATTGTGCAAGA

0                      410                      420                      430                      440                      450

Y D Y Y G S S Y F D Y W G Q G T T L T  
TACGATTACTACGGTAGTAGCTACTTTGACTACTGGGGCCAAGGGACCACGGTCACC  
CDR3                      BstEII

460                      470                      480                      490                      500                      510

V S S G G G G S G G G G S G G G G S Q  
GTCTCCTCAGGTGGAGGCGGTTTCAGGCGGAGGTGGCTCTGGCGGTGGCGGATCCCAG  
/------(G4S)<sub>3</sub> linker-----BamHI-/Vλ

520                      530                      540                      550                      560                      570

V V L T Q E S A L T T S P G E T V T L  
GCTGTTGTGACTCAGGAATCTGCACTCACCACATCACCTGGTGAAACAGTCACACTC

580                      590                      600                      610                      620

T C R S S T G A V T T S N Y A N W V Q  
ACTTGTCGCTCAAGTACTGGGGCTGTTACAAGTAACTATGCCAACTGGGTCCAA  
CDR1

FIGURE 4 (START)

10

630            640            650            660            670            680  
E K P D H L F T G L I G G T N N R A P  
GAAAAACCAGATCATTTATTCACTGGTCTAATAGGTGGTACCAACAACCGAGCTCCA  
*KpnI* CDR2 *SstI*

690 700 710 720 730 740  
G V P A R F S G S L I G D K A A L T I  
GGTGTTCCTGCCAGATTCTCAGGCTCCCTGATTGGAGACAAGGCTGCCCTCACCATC

750 760 770 780 790 8  
T G A Q T E D E A I Y F C A L W Y S N  
ACAGGGGCACAGACTGAGGATGAGGCAATATATTTCTGTGCTCTATGGTACAGCAAC  
CDR3

00           810                 820                 830                 840                 850  
H W V F G G G T K L T V L G L E \* \*  
CACTGGGTGTTCCGGTGAGGAACCAAACTGACTGTCCTAGGTCTCGAGTAATAAGA  
XhoI Eco

TTC  
RI

FIGURE 4 (END)




Fig. 5



Concentration of  
competing hapten  
(micrograms/ml)




Fig. 6




FIGURE 7




FIGURE 8

10/38

```

10      20      30      40      50      60
      M K Y L L P T
AAGCTTGCATGCAAATTCTATTTCAAGGAGACAGTCATAATGAAATACCTATTGCCTACG
HindIII SphI      SD      /-----

70      80      90      100      110      120
A A A G L L L L A A Q P A M A Q V Q L Q
GCAGCCGCTGGATTGTTATTACTCGCTGCCCAACCAGCGATGGCCAGGTGCAGCTGCAG
-----pelB leader-----/VH1NP PstI

130      140      150      160      170      180
Q P G A E L V K P G A S V K L S C K A S
CAGCCTGGGGCTGAGCTTGTGAAGCCTGGGGCTTCAGTGAAGCTGCTCTGCAAGGCTTCT

190      200      210      220      230      240
G Y T F T S Y W M H W V K Q R P G R G L
GGCTACACCTTCACCAGCTACTGGATGCACTGGGTGAAGCAGAGGCTGGACGAGGCCTT
CDR1

250      260      270      280      290      300
E W I G R I D P N S G G T R Y N E K F L
GAGTGGATTGGAAGGATTGATCCTAATAGTGGTGGTACTAAGTACATGAGAAGTTCAAG
CDR2

310      320      330      340      350      360
S K A T L T V D K P S S T A Y M Q L S S
AGCAAGGCCACACTGACTGTAGACAAACCCCTCCAGCACAGCCTACATGCAGCTCAGCAGC

370      380      390      400      410      420
L T S E D S A V Y Y C A R Y D Y Y G S S
CTGACATCTGAGGACTCTGCGGTCTATTATTGTGCAAGATACGATTACTACGGTAGTAGC

430      440      450      460      470      480
Y F D Y W G G Q G T T L T V S S G G G G S
TACTTTGACTACTGGGGCCAAGGGACCACGGTCACCGTCTCCTCAGGTGGAGGCGGTTCA
CDR3      BstEII      /---(G4S)3-----

490      500      510      520      530      540
G G G G S G G G G S Q V V L T Q E S A L
GGCGGAGGTGGCTCTGGCGGTGGCGGATCCAGGCTGTTGTGACTCAGGAATCTGCACTC
-----BamHI- /Vλ3558L

550      560      570      580      590      600
T T S P G E T V T L T C R S S T G A V T
ACCACATCACCTGGTGAAACAGTCACACTCACTTGTCTGCTCAAGTACTGGGGCTGTTACA
CDR1

610      620      630      640      650      660
T S N Y A N W V Q E K P D H L F T G L I
ACTAGTAACTATGCCAACTGGGTCCAAGAAAAACCAGATCATTTATCACTGGTCTAATA

670      680      690      700      710      720
G G T N N R A P G V P A R F S G S L I G
GGTGGTACCAACAACCGAGCTCCAGGTGTTCTCTGCCAGATTCTCAGGCTCCCTGATTGGA
KpnI CDR2 SstI

730      740      750      760      770      780
D K A A L T I T G A Q T E D E A I Y F C
GACAAGGCTGCCCTCACCATCACAGGGGCACAGACTGAGGATGAGGCAATATATTTCTGT

```

FIGURE 9 (START)

11/38

```

      790      800      810      820      830      840
A L W Y S N H W V F G G G T K L T V L G
GCTCTATGGGTACAGCAACCACTGGGTGTTCGGTGGAGGAACCAAAGTACTGTCCTAGGT
CDR3

      850      860      870      880      890      900
L E A P A A A P A D P S K D S K A Q V S
CTCGAGGCACCTGCTGCCGCACCTGCAGACCCGTCCAAGGACTCCAAAGCTCAGGTTTCT
XhoI /-Ala.Pro linker-----/Streptavidin Ps

      910      920      930      940      950      960
A A E A G I T G T W Y N Q L G S T F I V
GCAGCCGAAGCTGGTATCACTGGCACCTGGTATAACCAACTGGGGTTCGACTTTTCATTGTG
tI Sall

      970      980      990      1000      1010      1020
T A G A D G A L T G T Y E S A V G N A E
ACCGCTGGTGGGACGGAGCTCTGACTGGCACCTACGAATCTGCGGTTGGTAACGCAGAA

      1030      1040      1050      1060      1070      1080
S R Y V L T G R Y D S A P A T D G S G T
TCCCGCTACGTACTGACTGGCCGTTATGACTCTGCACCTGCCACCGATGGCTCTGGTACC
KpnI

      1090      1100      1110      1120      1130      1140
A L G W T V A W K N N Y R N A H S A T T
GCTCTGGGCTGGACTGTGGCTTGGAACCAACTATCGTAATGCGCACAGCGCCACTACG

      1150      1160      1170      1180      1190      1200
W S G Q Y V G G A E A R I N T Q W L L T
TGGTCTGGCCAATACGTTGGCGGTGCTGAGGCTCGTATCAACACTCAGTGGCTGTAAACA

      1210      1220      1230      1240      1250      1260
S G T T E A N A W K S T L V G H D T F T
TCCGGCACTACCGAAGCGAATGCATGGAAATCGACACTAGTAGGTCATGACACCTTTACC

      1270      1280      1290      1300      1310      1320
K V K P S A A S I D A A K K A G V N N G
AAAGTTAAGCCTTCTGCTGCTAGCATTGATGCTGCCAAGAAAGCAGGCGTAAACAACGGT
Bst

      1330      1340      1350      1360
N P L D A V Q Q * *
AACCTCTAGACGCTGTTTCAGCAATAATAAGAATT
EII XbaI EcoRI

```

FIGURE 9 (END)

12/38

```

      10          20          30          40          50          60
      M K Y L L P T
AAGCTTGCATGCAAATTCTATTTCAGGAGACAGTCATAATGAAATACCTATTGCCTACG
HindIIISphI                      SD /-----

      70          80          90          100         110         120
A A A G L L L L A A Q P A M A Q V Q L Q
GCAGCCGCTGGATTGTTATTACTCGCTGCCCAACCAGCGATGGCCAGGTGCAGCTGCAG
-----pelB leader-----/VHNP PstI

      130         140         150         160         170         180
Q P G A E L V K P G A S V K L S C K A S
CAGCCTGGGGCTGAGCTTGTGAAGCCTGGGGCTTCAGTGAAGCTGTCCTGCAAGGCTTCT

      190         200         210         220         230         240
G Y T F T S Y W M H W V K Q R P G R G L
GGCTACACCTTCACCACTACTGGATGCACTGGGTGAAGCAGAGGCCTGGACGAGGCCTT
      CDR1

      250         260         270         280         290         300
E W I G R I D P N S G G T R Y N E K F L
GAGTGGATTGGAAGGATTGATCCTAATAGTGGTGGTACTAAGTACAATGAGAAGTTCAAG
      CDR2

      310         320         330         340         350         360
S K A T L T V D K P S S T A Y M Q L S S
AGCAAGGCCACACTGACTGTAGACAAACCCTCCAGCACAGCCTACATGCAGCTCAGCAGC

      370         380         390         400         410         420
L T S E D S A V Y Y C A R Y D Y Y G S S
CTGACATCTGAGGACTCTGCGGTCTATTATTGTGCAAGATACGATTACTACGGTAGTAGC

      430         440         450         460         470         480
Y F D Y W G Q G T T L T V S S G G G G S
TACTTTGACTACTGGGGCCAAGGGACCACGGTCAACCGTCTCCTCAGGTGGAGGCGGTTCA
CDR3                      BstEII /---(G4S)3-----

      490         500         510         520         530         540
G G G G S G G G G S Q V V L T Q E S A L
GGCGGAGGTGGCTCTGGCGGTGGCGGATCCAGGCTGTTGTGACTCAGGAATCTGCACTC
-----BamHI-/VλJ558L

      550         560         570         580         590         600
T T S P G E T V T L T C R S S T G A V T
ACCACATCACCTGGTGAAACAGTCACACTCACTGTCTCGCTCAAGTACTGGGGCTGTTACA
      CDR1

      610         620         630         640         650         660
T S N Y A N W V Q E K P D H L F T G L I
ACTAGTAAGTATGCCAAGTGGGTCCAAGAAAAACCAGATCATTATTCACTGGTCTAATA

      670         680         690         700         710         720
G G T N N R A P G V P A R F S G S L I G
GGTGGTACCAACAACCGAGCTCCAGGTGTTCTCCTGCCAGATTCTCAGGCTCCCTGATTGGA
KpnI CDR2 SstI

      730         740         750         760         770         780
D K A A L T I T G A Q T E D E A I Y F C
GACAAGGCTGCCCTCACCATCACAGGGGCACAGACTGAGGATGAGGCAATATATTTCTGT

```

FIGURE 10 (START)

13/38

```

      790      800      810      820      830      840
A L W Y S N H W V F G G G T K L T V L G
GCTCTATGGTACAGCAACCACCTGGGTGTTCGGTGGAGGAACCAAAGTACTGTCCTAGGT
CDR3

      850      860      870      880      890      900
L E A P A A A P A D P S K D S K A Q V S
CTCGAGGCACCTGCTGCCGCACCTGCAGACCCGTCCAAGGACTCCAAAGCTCAGGTTTCT
XhoI /-Ala.Pro linker-----/Streptavidin Ps

      910      920      930      940      950      960
A A E A G I T G T W Y N Q L G S T F I V
GCAGCCGAAGCTGGTATCACTGGCACCTGGTATAACCAACTGGGGTTCGACTTTTCATTGTG
EI Sall

      970      980      990      1000      1010      1020
T A G A D G A L T G T Y E S A V G N A E
ACCGCTGGTGGCGGACGGAGCTCTGACTGGCACCTACGAATCTGCGGTTGGTAACGCAGAA

      1030      1040      1050      1060      1070      1080
S R Y V L T G R Y D S A P A T D G S G T
TCCCGCTACGTACTGACTGGCCGTTATGACTCTGCACCTGCCACCGATGGCTCTGGTACC
KpnI

      1090      1100      1110      1120      1130      1140
A L G W T V A W K N N Y R N A H S A T T
GCTCTGGGCTGGACTGTGGCTTGAAAAACAATATCGTAATGCGCACAGCGCCACTACG

      1150      1160      1170      1180      1190      1200
W S G Q Y V G G A E A R I N T Q W L L T
TGGTCTGGCCAATACGTTGGCGGTGCTGAGGCTCGTATCAACACTCAGTGGCTGTTAACA

      1210      1220      1230      1240      1250      1260
S G T T E A N A W K S T L V G H D T F T
TCCGGCACTACCGAAGCGAATGCATGGAAATCGACACTAGTAGGTCATGACACCTTTACC

      1270      1280      1290
K V K P S A A S * *
AAAGTTAAGCCTTCTGCTGCTAGCTAATAAGAAATC
EcoRI

```

FIGURE 10 (END)

14/38

```

10      20      30      40      50      60
AAGCTTGCATGCAAATTCTATTTCAAGGAGACAGTCATAATGAAATACCTATTGCCTACG
HindIIISphI                      SD /-----

70      80      90      100     110     120
A A A G L L L L A A Q P A M A Q V Q L Q
GCAGCCGCTGGATTGTTATTACTCGCTGCCCAACCAGCGATGSCCAGGTGCAGCTGCAG
-----pelB leader-----/VHNP PstI

130     140     150     160     170     180
Q P G A E L V K P G A S V K L S C K A S
CAGCCTGGGGCTGAGCTTGTGAAGCCTGGGGCTTCAGTGAAGCTGTCTGCAAGGCTTCT

190     200     210     220     230     240
G Y T F T S Y W M H W V K Q R P G R G L
GGCTACACCTTCACCAGCTACTGGATGCACTGGGTGAAGCAGAGGCCTGGACGAGGCCTT
CDR1

250     260     270     280     290     300
E W I G R I D P N S G G T R Y N E K F L
GAGTGGATTGGAAGGATTGATCCTAATAGTGGTGGTACTAAGTACAATGAGAAGTTCAAG
CDR2

310     320     330     340     350     360
S K A T L T V D K P S S T A Y M Q L S S
AGCAAGGCCACACTGACTGTAGACAAACCTCCAGCACAGCCTACATGCAGCTCAGCAGC

370     380     390     400     410     420
L T S E D S A V Y Y C A R Y D Y Y G S S
CTGACATCTGAGGACTCTGCGGTCTATTATTGTGCAAGATACGATTACTACGGTAGTAGC

430     440     450     460     470     480
Y F D Y W G Q G T T L T V S S G G G G S
TACTTTGACTACTGGGGCCAAGGGACCCAGGTCACCGTCTCCTCAGGTGGAGGCGGTTCA
CDR3                      BstEII /---(G4S)3-----

490     500     510     520     530     540
G G G G S G G G G S Q V V L T Q E S A L
GGCGGAGGTGGCTCTGGCGGTGGCGGATCCAGGCTGTTGTGACTCAGGAATCTGCACTC
-----BamHI-/VλJ558L

550     560     570     580     590     600
T T S P G E T V T L T C R S S T G A V T
ACCACATCACCTGGTGAAACAGTCACACTCACTTGTGCGCTCAAGTACTGGGGCTGTTACA
CDR1

610     620     630     640     650     660
T S N Y A N W V Q E K P D H L F T G L I
ACTAGTAACTATGCCAACTGGGTCCAAGAAAAACCAGATCATTTATTCACTGGTCTAATA

670     680     690     700     710     720
G G T N N R A P G V P A R F S G S L I G
GGTGGTACCAACAACCGAGCTCCAGGTGTTCTGCTGCCAGATTCTCAGGCTCCCTGATTGGA
KpnI CDR2 SstI

730     740     750     760     770     780
D K A A L T I T G A Q T E D E A I Y F C
GACAAGGCTGCCCTCACCATCACAGGGGCACAGACTGAGGATGAGGCAATATATTTCTGT

```

FIGURE 11 (START)



790            800            810            820            830            840  
 A L W Y S N H W V F G G G T K L T V L G  
GCTCTATGGTACAGCAACCACCTGGGTGTTTCGGTGGAGGAACCAACTGACTGTCCTAGGT  
 CDR3

850            860            870            880            890            900  
 L E A P A A A P A E A G I T G T W Y N Q  
CTCGAGGCACCTGCTGCCGCACCTGCCGAAGCTGGTATCACTGGCACCTGGTATAACCAA  
 XhoI /-Ala.Pro linker--/core streptavidin

910            920            930            940            950            960  
 L G S T F I V T A G A D G A L T G T Y E  
 CTGGGGTCGACTTTCATTGTGACCGCTGCTGCGGACGGAGCTCTGACTGGCACCTACGAA  
 SalI

970            980            990            1000            1010            1020  
 S A V G N A E S R Y V L T G R Y D S A P  
 TCTGCGGTGGTAACGCAGAATCCCGCTACGTACTGACTGGCCGTTATGACTCTGCACCT

1030            1040            1050            1060            1070            1080  
 A T D G S G T A L G W T V A W K N N Y R  
 GCCACCGATGGCTCTGGTACCGCTCTGGGCTGGACTGTGGCTTGAAAAACAACTATCGT  
 KpnI

1090            1100            1110            1120            1130            1140  
 N A H S A T T W S G Q Y V G G A E A R I  
 AATGCGCACAGCGCCACTACGTGGTCTGGCCAATACGTTGGCGGTGCTGAGGCTCGTATC

1150            1160            1170            1180            1190            1200  
 N T Q W L L T S G T T E A N A W K S T L  
 AACACTCAGTGGCTGTAAACATCCGGCACTACCGAAGCGAATGCATGGAAATCGACACTA

1210            1220            1230            1240            1250            1260  
 V G H D T F T K V K P S A A S \* \*  
 GTAGGTCATGACACCTTTACCAAAGTTAAGCCTTCTGCTGCTAGCTAATAAGAATTC  
 EcoRI

FIGURE 11 (END)




Fig. 12




Fig. 13





Fig. 14





FIGURE 15







FIGURE 17








Fig.19




Fig. 20

AAGCTTGCATGCAAATTCTATTTCAAGGAGACAGTCATAATGAAATACCTA  
TTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCCAACCAGCGATG  
GCCCAGGTGCAGCTGCAGCAGCCTGGGGCTGAGCTTGTGAAGCCTGGGGCT  
TCAGTGAAGCTGTCCTGCAAGGCTTCTGGCTACACCTTCACCAGCTACTGG  
ATGCACTGGGTGAAGCAGAGGCCTGGACGAGGCCTTGAGTGGATTGGAAGG  
ATTGATCCTAATAGTGGTGGTACTAAGTACAATGAGAAGTTCAAGAGCAAG  
GCCACACTGACTGTAGACAAACCCTCCAGCACAGCCTACATGCAGCTCAGC  
AGCCTGACATCTGAGGACTCTGCGGTCTATTATTGTGCAAGATACGATTAC  
TACGGTAGTAGCTACTTTGACTACTGGGGCCAAGGGACCACGGTCACCGTC  
TCCTCAGGTGGAGGCGGTTTCAGGCGGAGGTGGCTCTGGCGGTGGCGGATCC  
CAGGCTGTTGTGACTCAGGAATCTGCACTCACCACATCACCTGGTGAAACA  
GTCACACTCACTTGTCTGCTCAAGTACTGGGGCTGTTACAAGTAGTAAGTAT  
GCCAACTGGGTCCAAGAAAAACCAGATCATTTATTCACTGGTCTAATAGGT  
GGTACCAACAACCGAGCTCCAGGTGTTCTGCCAGATTCTCAGGCTCCCTG  
ATTGGAGACAAGGCTGCCCTCACCATCACAGGGGCACAGACTGAGGATGAG  
GCAATATATTTCTGTGCTCTATGGTACAGCAACCACTGGGTGTTTCGGTGGA  
GGAACCAAACTGACTGTCCTAGGTCTCGAGATCAAGCGCAAGGAATCTGCA  
GCTGCCAAGTTCGAGCGGCAGCACATGGACTCTGGCAACTCCCCCAGCAGC  
AGCTCCAACCTACTGCAACCTGATGATGTGCTGCCGAAGATGACCCAGGGGA  
AATGCAAGCCAGTGAACACCTTTGTGCATGAGTCCCTGGCCGATGTTAAGG  
CCGTGTGCTCCCAGAAGAAAGTCACTTGCAAGAATGGGCAGACCAACTGCT  
ACCAGAGCAAATCCACCATGCGCATCACAGACTGCCGCGAGACTGGCAGCT  
CCAAGTACCCCAACTGCGCCTACAAGACCACCCAGGTGGAGAAACACATCA  
TAGTGGCTTGTGGCGGTAAACCGTCCGTGCCAGTCCACTTCGATGCTTCAG  
TG TAGATCTCCACCTGAGGCCAGAACAGTGAATTC

FIGURE 21

AAGCTTGCATGCAAATTCTATTTCAAGGAGACAGTCATAATGAAATACCTA  
TTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCAACCAGCGATG  
GCCCAGGTGCAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAGAGCCTG  
TCCATCACATGCACTGTCTCAGGGTTCTCATTAAACCAGTTATGGTGTAAAGC  
TGGGTTCGCCAGCCTCCAAGAAAGGGTCTGGAGTGGCTGGGAGTAATATGG  
GAAGACGGGAGCACAAATTATCATTACGTCTCATATCCAGACTGAGCATC  
AACAAGGATAACTCCAAGAGCCAAGTTTTCTTAAACTGAACAGTCTGCAA  
ACTGATGACACAGCCACGTACTACTGTGCCAAACCCCACTACGGTAGCAGC  
AACGTGGGGGCTATGGAATACTGGGGTCAAGGAACCTCGGTCACCGTCTCC  
TCAGGTGGAGGCGGTTTCAGGCGGAGGTGGCTCTGGCGGTGGCGGATCGGAC  
ATCGAGCTCACCCAGTCTCCAGCCTCCCTAACTGCATCTGTGGGAGAACT  
GTCACCATCACCTGTCGAGCAAGTGAAAATATTTACAGTTATGTAGCATGG  
TATCAGCAGAAACAGGGAAAATCTCCTCAGTTCCTGGTCTATAATGCAAAA  
TCCTTAGCAGAGGGTGTGCCATCAAGGTTTCAGTGGCAGTGGATCAGGCACA  
CAGTTTTCTCTGAAGATCAACAGCCTGCAGCCTGAAAATTTTGGGAATTAT  
TACTGTCAACATCATTATGTTAGTCCGTGGACGTTTCGGTGGAGGCACCAAG  
CTCGAGATCAAGCGCAAGGAATCTGCAGCTGCCAAGTTCGAGCGGCAGCAC  
ATGGACTCTGGCAACTCCCCCAGCAGCAGCTCCAATACTGCAACCTGATG  
ATGTGCTGCCGAAGATGACCCAGGGGAAATGCAAGCCAGTGAACACCTTTG  
TGCATGAGTCCCTGGCCGATGTTAAGGCCGTGTGCTCCCAGAAGAAAGTCA  
CTTGCAAGAATGGGCAGACCAACTGCTACCAGAGCAAATCCACCATGCGCA  
TCACAGACTGCCGCGAGACTGGCAGCTCCAAGTACCCCAACTGCGCCTACA  
AGACCACCCAGGTGGAGAAACACATCATAGTGGCTTGTGGCGGTAAACCGT  
CCGTGCCAGTCCACTTCGATGCTTCAGTGTAGATCTCCACCTGAGGCCAGA  
ACAGTGAATTC

AAGCTTG CATGCAAATTCTATTTCAAGGAGACAGTCATAATGAAATACCTA  
TTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCCAACCAGCGATG  
GCCCAGGTGCAGCTGCAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAG  
ACGCTGTCCATCACATGCACCGTCTCAGGGTTCTCATTAACCGGCTATGGT  
GTAAACTGGGTTTCGCCAGCCTCCAGGAAAGGGTCTGGAGTGGCTGGGAATG  
ATTTGGGGTGATGGAAACACAGACTATAATT CAGCTCTCAAATCCAGACTG  
AGCATCAGCAAGGACAAC TCCAAGAGCCAAGTTTTCTTAAAAATGAACAGT  
CTGCACACTGATGACACAGCCAGG TACTACTGTGCCAGAGAGAGAGATTAT  
AGGCTTGACTACTGGGGCCAAGGCACCACGGTCACCGTCTCCTCAGGTGGA  
GGCGGTT CAGGCGGAGGTGGCTCTGGCGGTGGCGGATCGGACATCGTCATG  
ACTCAGTCTCCAGCCTCCCTTTCTGCGTCTGTGGGAGAAACTGTCACCATC  
ACATGTCGAGCAAGTGGGAATATTCACAATTATTTAGCATGGTATCAGCAG  
AAACAGGGGAAAATCTCCTCAGCTCCTGGTCTATTATACAACAACCTTAGCA  
GATGGTGTGCCATCAAGGTT CAGTGGCAGTGGATCAGGAACACAATATTCT  
CTCAAGATCAACAGCCTGCAGCCTGAAGATTTTGGGAGTTATTACTGTCAA  
CATTTTTTGGAGTACTCCTCGGACGTTCCGGTGGAGGCACCAAGCTCGAGATC  
AAGCGCAAGGAATCTGCAGCTGCCAAGTTCGAGCGGCAGCACATGGACTCT  
GGCAACTCCCCCAGCAGCAGCTCCA ACTACTGCAACCTGATGATGTGCTGC  
CGAAGATGACCCAGGGGAAATGCAAGCCAGTGAACACCTTTGTGCATGAGT  
CCCTGGCCGATGTTAAGGCCGTGTGCTCC CAGAAGAAAGTCACTTGCAAGA  
ATGGGCAGACCAACTGCTACCAGAGCAAATCCACCATGCGCATCACAGACT  
GCCGCGAGACTGGCAGCTCCAAGTACCCCAACTGCGCCTACAAGACCACCC  
AGGTGGAGAAACACATCATAGTGGCTTGTGGCGGTAAACCGTCCGTGCCAG  
TCCACTTCGATGCTTCAGTGTAGATCTCCACCTGAGGCCAGAACAGTGAAT  
TC

FIGURE 23

AAGCTTGCATGCAAATTCTATTTCAAGGAGACAGTCATAATGAAATACCTA  
TTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCCAACCAGCGATG  
GCCCAGGTGCAGCTGCAGCAGCCTGGGGCTGAGCTTGTGAAGCCTGGGGCT  
TCAGTGAAGCTGTCCTGCAAGGCTTCTGGCTACACCTTCACCAGCTACTGG  
ATGCACTGGGTGAAGCAGAGGCCTGGACGAGGCCTTGAGTGGATTGGAAGG  
ATTGATCCTAATAGTGGTGGTACTAAGTACAATGAGAAGTTCAAGAGCAAG  
GCCACACTGACTGTAGACAAACCCTCCAGCACAGCCTACATGCAGCTCAGC  
AGCCTGACATCTGAGGACTCTGCGGTCTATTATTGTGCAAGATACGATTAC  
TACGGTAGTAGCTACTTTGACTACTGGGGCCAAGGGACCACGGTCACCGTC  
TCCTCAGGTGGAGGCGGTTCCAGGCGGAGGTGGCTCTGGCGGTGGCGGATCC  
CAGGCTGTTGTGACTCAGGAATCTGCACTCACCACATCACCTGGTGAAACA  
GTCACACTCACTTGTGCTCAAGTACTGGGGCTGTTACAACCTAGTAACTAT  
GCCAACTGGGTCCAAGAAAAACCAGATCATTATTTACTGGTCTAATAGGT  
GGTACCAACAACCGAGCTCCAGGTGTTCCCTGCCAGATTCTCAGGCTCCCTG  
ATTGGAGACAAGGCTGCCCTCACCATCACAGGGGCACAGACTGAGGATGAG  
GCAATATATTTCTGTGCTCTATGGTACAGCAACCACTGGGTGTTCCGGTGGA  
GGAACCAAACTGACTGTCCTAGGTCTCGAGATTAAACGTATGCTTAAGATC  
GCTGCTTTCAACATACGTACCTTCGGTGAATCTAAAATGTCTAACGCTACG  
CTAGCATCTTACATCGTACGCATCGTACGCCGTTACGATATCGTTCTGATC  
CAGGAAGTTCGCGACTCTCACCTGGTTGCAGTTGGTAACTTCTAGACTAC  
CTGAACCAGGACGACCCGAACACCTACCACTACGTTGTTTCTGAACCCCTC  
GGGCGTAACTCTTACAAAGAACGGTACCTGTTCCCTGTTCCGTCCGAACAAA  
GTTTCAGTACTGGATACCTACCAGTACGACGACGGATGCGAATCTTGCGGT  
AACGACTCTTTCTCCCGGGAACCGGCTGTTGTTAAATTCTCGAGCCACTCT  
ACCAAGGTAAAGAGTTCGCTATCGTTGCTCTGCACAGCGCGCGCTCTGAC  
GCTGTTGCTGAAATCAACTCTCTGTACGACGTTTACCTGGACGTTTCAGCAG  
AAATGGCACCTGAACGACGTCATGCTGATGGGTGACTTCAACGCTGACTGC  
TCTTATGTAACCTCTTCTCAGTGGTCATCGATTTCGTCTGCGCACCTCGTCC  
ACCTTCCAGTGGCTGATCCCGGACTCCGCTGACACCACCGCTACTAGTACC  
AACTGCGCTTACGACCGTATCGTTGTTGCTGGATCCCTGCTGCAGTCTTCT  
GTTGTACCGGGTAGCGCGGCCCGTTTCGACTTCCAGGCTGCATATGGTCTT  
TCGAACGAAATGGCGCTGGCCATCTCTGATCACTACCCGGTTGAGGTAACC  
CTGACCTAATTCTAGA

AAGCTTGCATGCAAATTCTATTTCAAGGAGACAGTCATAATGAAATACCTA  
TTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCAACCAGCGATG  
GCCCAGGTGCAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAGAGCCTG  
TCCATCACATGCACTGTCTCAGGGTTCTCATTAACCAGTTATGGTGTAAGC  
TGGGTTCGCCAGCCTCCAAGAAAGGGTCTGGAGTGGCTGGGAGTAATATGG  
GAAGACGGGAGCACAAATTATCATTACGTCTCATATCCAGACTGAGCATC  
AACAAGGATAACTCCAAGAGCCAAGTTTTCTTAAACTGAACAGTCTGCAA  
ACTGATGACACAGCCACGTACTACTGTGCCAAACCCCACTACGGTAGCAGC  
AACGTGGGGGCTATGGAATACTGGGGTCAAGGAACCTCGGTACCGTCTCC  
TCAGGTGGAGGCGGTTTCAGGCGGAGGTGGCTCTGGCGGTGGCGGATCGGAC  
ATCGAGCTACCCAGTCTCCAGCCTCCCTAACTGCATCTGTGGGAGAAACT  
GTCACCATCACCTGTCGAGCAAGTGAAAATATTTACAGTTATGTAGCATGG  
TATCAGCAGAAACAGGGGAAAATCTCCTCAGTTCCTGGTCTATAATGCAAAA  
TCCTTAGCAGAGGGTGTGCCATCAAGGTTTCAGTGGCAGTGGATCAGGCACA  
CAGTTTTCTCTGAAGATCAACAGCCTGCAGCCTGAAAATTTTGGAATTAT  
TACTGTCAACATCATTATGTTAGTCCGTGGACGTTTCGGTGGAGGCACCAAG  
CTCGAGATTAAACGTATGCTTAAGATCGCTGCTTTCAACATACGTACCTTC  
GGTGAATCTAAAATGTCTAACGCTACGCTAGCATCTTACATCGTACGCATC  
GTACGCCGTTACGATATCGTTCTGATCCAGGAAGTTCGCGACTCTCACCTG  
GTTGCAGTTGGTAAACTTCTAGACTACCTGAACCAGGACGACCCGAACACC  
TACCACTACGTTGTTTCTGAACCCCTCGGGCGTAACTCTTACAAAGAACGG  
TACCTGTTCCCTGTTCCGTCCGAACAAAGTTTCAGTACTGGATACCTACCAG  
TACGACGACGGATGCGAATCTTGCGGTAACGACTCTTTCTCCCGGGAACCG  
GCTGTTGTTAAATTCTCGAGCCACTCTACCAAGGTAAAGAGTTCGCTATC  
GTTGCTCTGCACAGCGCGCCGTCTGACGCTGTTGCTGAAATCAACTCTCTG  
TACGACGTTTACCTGGACGTTTCAGCAGAAATGGCACCTGAACGACGTCATG  
CTGATGGGTGACTTCAACGCTGACTGCTCTTATGTAACCTCTTCTCAGTGG  
TCATCGATTTCGTCTGCGCACCTCGTCGACCTTCCAGTGGCTGATCCCGGAC  
TCCGCTGACACCACCGCTACTAGTACCAACTGCGCTTACGACCGTATCGTT  
GTTGCTGGATCCCTGCTGCAGTCTTCTGTTGTACCGGGTAGCGCGGCCCCG  
TTCGACTTCCAGGCTGCATATGGTCTTTCGAACGAAATGGCGCTGGCCATC  
TCTGATCACTACCCGGTTGAGGTAACCCTGACCTAATTCTAGA

FIGURE 25

AAGCTTG CATGCAAATTCTATTTCAAGGAGACAGTCATAATGAAATACCTA  
TTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCAACCAGCGATG  
GCCCAGGTGCAGCTGCAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAG  
ACGCTGTCCATCACATGCACCGTCTCAGGGTTCTCATTAAACCGGCTATGGT  
GTAAACTGGGTTTCGCCAGCCTCCAGGAAAGGGTCTGGAGTGGCTGGGAATG  
ATTTGGGGTGATGGAAACACAGACTATAATTCAGCTCTCAAATCCAGACTG  
AGCATCAGCAAGGACAACCTCCAAGAGCCAAGTTTTCTTAAAAATGAACAGT  
CTGCACACTGATGACACAGCCAGGTACTACTGTGCCAGAGAGAGAGATTAT  
AGGCTTGACTACTGGGGCCAAGGCACCAACGGTCACCGTCTCCTCAGGTGGA  
GGCGGTTTCAGGCGGAGGTGGCTCTGGCGGTGGCGGATCGGACATCGTCATG  
ACTCAGTCTCCAGCCTCCCTTTCTGCGTCTGTGGGAGAACTGTCACCATC  
ACATGTGCGAGCAAGTGGGAATATTACAAATTATTTAGCATGSTATCAGCAG  
AAACAGGGAAAATCTCCTCAGCTCCTGGTCTATTATACAACAACCTTAGCA  
GATGGTGTGCCATCAAGGTTCAGTGGCAGTGGATCAGGAACACAATATTCT  
CTCAAGATCAACAGCCTGCAGCCTGAAGATTTTGGGAGTTATTACTGTCAA  
CATTTTTGGAGTACTCCTCGGACGTTTCGGTGGAGGCACCAAGCTCGAGATT  
AAACGTATGCTTAAGATCGCTGCTTTCAACATACGTACCTTCGGTGAATCT  
AAAATGTCTAACGCTACGCTAGCATCTTACATCGTACGCATCGTACGCCGT  
TACGATATCGTTCTGATCCAGGAAGTTCGCGACTCTCACCTGGTTGCAGTT  
GGTAAACTTCTAGACTACCTGAACCAGGACGACCCGAACACCTACCACTAC  
GTTGTTTCTGAACCCCTCGGGCGTAACTCTTACAAAGAACGGTACCTGTTCT  
CTGTTCCGTCCGAACAAAGTTTCAGTACTGGATACCTACCAGTACGACGAC  
GGATGCGAATCTTGCGGTAAACGACTCTTTCTCCCGGGAACCGGCTGTTGTT  
AAATTCTCGAGCCACTCTACCAAGGTTAAAGAGTTCGCTATCGTTGCTCTG  
CACAGCGCGCCGTCTGACGCTGTTGCTGAAATCAACTCTCTGTACGACGTT  
TACCTGGACGTTTACGAGAAATGGCACCTGAACGACGTCATGCTGATGGGT  
GACTTCAACGCTGACTGCTCTTATGTAACCTCTTCTCAGTGGTCATCGATT  
CGTCTGCGCACCTCGTCGACCTTCCAGTGGCTGATCCCGGACTCCGCTGAC  
ACCACCGCTACTAGTACCAACTGCGCTTACGACCGTATCGTTGTTGCTGGA  
TCCCTGCTGCAGTCTTCTGTTGTACCGGGTAGCGCGGCCCCGTTCGACTTC  
CAGGCTGCATATGGTCTTTCCGAACGAAATGGCGCTGGCCATCTCTGATCAC  
TACCCGGTTGAGGTAACCCTGACCTAATTCTAGA

FIGURE 26






Fig. 27




Fig. 28

ATGAAATACCTATTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCCAACC  
AGCGATGGCCCAGCTGCAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAGAGCC  
TGTCATCACATGCACTGTCTCAGGGTTCTCATTAAACCAGTTATGGTGTAAGCTGG  
GTTCCGCCAGCCTCCAACTCAGGGTCTGGAGTGGCTGGGAGTAATATGGGAAGACGG  
GAGCACAAATCTCTCATATCCAGACTGAGCATCAACAAGGATAACT  
CCAAGAGCCAACTCTTAAACTGAACAGTCTGCAAACTGATGACACAGCCACG  
TACTACTGTGCCAAACCCCACTACGGTAGCAGCAACGTGGGGGCTATGGAATACTG  
GGGTCAAGGAACCTCGGTCACCGTCTCCTCAGGTGGAGGCGGTTCAGGCGGAGGTG  
GCTCTGGCGGTGGCGGATCGGACATCGAGCTCACCCAGTCTCCAGCCTCCCTAACT  
GCATCTGTGGGAGAACTGTCACCATCACCTGTGAGCAAGTGAAAATATTTACAG  
TTATGTAGCATGGTATCAGCAGAAACAGGGAAAATCTCCTCAGTTCCTGGTCTATA  
ATGCAAAATCCTTAGCAGAGGGTGTGCCATCAAGGTTCAAGTGGCAGTGGATCAGGC  
ACACAGTTTTCTCTGAAGATCAACAGCCTGCAGCCTGAAGATTTTGGGAATTATTA  
CTGTCAACATCATTATGTTAGTCCGTGGACGTTTCGGTGGAGGCACCAAGCTCGAGA  
TCAAGCGCTCTAGCCTCGAAGGTGGGTGCGCTGGTAATAGAGTCAGAAGATCAGTC  
GGAAGCAGCCTGTCTTGGGTGGTCTCGACGTCGAGATCAAGCGCAAGGAATCTGC  
AGCTGCCAAGTTCGAGCGGCAGCACATGGACTCTGGCAACTCCCCCAGCAGCAGCT  
CCAATACTGCAACCTGATGATGTGCTGCCGGAAGATGACCCAGGGGAAATGCAAG  
CCAGTGAACACCTTTGTGCATGAGTCCCTGGCCGATGTTAAGGCCGTGTGCTCCCA  
GAAGAAAGTCACTTGCAAGAATGGGCAGACCAACTGCTACCAGAGCAAATCCACCA  
TGCGCATCACAGACTGCCGCGAGACTGGCAGCTCCAAGTACCCCAACTGCGCCTAC  
AAGACCACCCAGGTGGAGAAACATCATAGTGGCTTGTGGCGGTAAACCGTCCGT  
GCCAGTCCACTTCGATGCTTCAGTGTAG

FIGURE 29

ATGAAATACCTATTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCAACC  
AGCGATGGCCCAGCTGCAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAGAGCC  
TGTCCATCACATGCACTGTCTCAGGGTTCTCATTAAACCAGTTATGGTGTAAAGCTGG  
GTTTCGCCAGCCTCCAAGAAAGGGTCTGGAGTGGCTGGGAGTAATATGGGAAGACGG  
GAGCACAATTATCATTCACGTCTCATATCCAGACTGAGCATCAACAAGGATAACT  
CCAAGAGCCAAGTTTTCTTAAACTGAACAGTCTGCAAACTGATGACACAGCCACG  
TACTACTGTGCCAAACCCACTACGGTAGCAGCAACGTGGGGGGCTATGGAATACTG  
GGGTCAAGGAACCTCGGTACCGTCTCCTCAGGTGGAGGCGGTTACAGGCGGAGGTG  
GCTCTGGCGGTGGCGGATCGGACATCGAGCTCACCCAGTCTCCAGCCTCCCTAACT  
GCATCTGTGGGAGAACTGTCACCATCACCTGTCGAGCAAGTGAAAATATTTACAG  
TTATGTAGCATGGTATCAGCAGAAACAGGGAAGATCTCCTCAGTTCCTGGTCTATA  
ATGCAAAATCCTTAGCAGAGGGTGTGCCATCAAGGTTCAAGTGGCAGTGGATCAGGC  
ACACAGTTTTCTCTGAAGATCAACAGCCTGCAGCCTGAAAATTTTGGGAATTATTA  
CTGTCAACATCATTATGTTAGTCCGTGGACGTTCCGTGGAGGCACCAAGCTCGAGA  
TCAAGCGCTCTAGCCTCGAAGGTGGGTGCGCTGGTAATAGAGTCAGAAGATCAGTC  
GGAAGCAGCCTGTCTTGCGGTGGTCTCGACGTCTGAGATCAAGCGCAAGGAATCTGC  
AGCTGCCAAGTTCGAGCGGCAGCACATGGACTCTGGCAACTCCCCAGCAGCAGCT  
CCAATACTGCAACCTGATGATGTGCTGCCGGAAGATGACCCAGGGGAAATGCAAG  
CCAGTGAACACCTTTGTGCATGAGTCCCTGGCCGATGTTAAGGCCGTGTGCTCCCA  
GAAGAAAGTCACTTGCAAGAATGGGCAGACCAACTGCTACCAGAGCAAATCCACCA  
TGCGCATCACAGACTGCCGCGAGACTGGCAGCTCCAAGTACCCCAACTGCGCCTAC  
AAGACCACCCAGGTGGAGAAACACATCATAGTGGCTTGTGGCGGTAAACCGTCCGT  
GCCAGTCCACTTCGATGCTTCAGTGAAGGACGAACTGTAA

ATGAAATACCTATTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCCAACC  
AGCGATGGCCCAGCTGCAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAGAGCC  
TGTCCATCACATGCACTGTCTCAGGGTTCTCATTAACCAGTTATGGTGTAAAGCTGG  
GTTCCGCCAGCCTCCAAGAAAGGGTCTGGAGTGGCTGGGAGTAATATGGGAAGACGG  
GAGCACAAATTATCATTACGTCTCATATCCAGACTGAGCATCAACAAGGATAACT  
CCAAGAGCCAAGTTTTCTTAAAACTGAACAGTCTGCAAACTGATGACACAGCCACG  
TACTACTGTGCCAAACCCCCACTACGGTAGCAGCAACGTGGGGGCTATGGAATACTG  
GGGTCAAGGAACCTCGGTACCGTCTCCTCAGGTGGAGGCGGTTTCAGGCGGAGGTG  
GCTCTGGCGGTGGCGGATCGGACATCGAGCTCACCCAGTCTCCAGCCTCCCTAACT  
GCATCTGTGGGAGAACTGTCACCATCACCTGTCTGAGCAAGTGAATAATTTTACAG  
TTATGTAGCATGGTATCAGCAGAAACAGGGAAAATCTCCTCAGTTCCTGGTCTATA  
ATGCAAAATCCTTAGCAGAGGGTGTGCCATCAAGGTTTCAGTGGCAGTGGATCAGGC  
ACACAGTTTTTCTCTGAAGATCAACAGCCTGCAGCCTGAAGATTTTGGGAATTATTA  
CTGTCAACATCATTATGTTAGTCCGTGGACGTTTCGGTGGAGGCACCAAGCTCGAGA  
TCAAGCGCTCTAGCCTCGAAGGTGGGTGCGCTGGTAATAGAGTCAGAAGATCAGTC  
GGAAGCAGCCTGTCTTGCGGTGGTCTCGACGTGAGATCAAGGCACCTGCTGCCTC  
CCCGGCAGACGCTAAGGAATCTGCAGCTGCCAAGTTCGAGCGGCAGCACATGGACT  
CTGGCAACTCCCCCAGCAGCAGCTCCAATACTGCAACCTGATGATGTGCTGCCGG  
AAGATGACCCAGGGGAAATGCAAGCCAGTGAACACCTTTGTGCATGAGTCCCTGGC  
CGATGTTAAGGCCGTGTGCTCCCAGAAGAAAGTCACTTGCAAGAATGGGCAGACCA  
ACTGCTACCAGAGCAAATCCACCATGCGCATCACAGACTGCCGCGAGACTGGCAGC  
TCCAAGTACCCCAACTGCGCCTACAAGACCACCCAGGTGGAGAAACACATCATAGT  
GGCTTGTTGGCGGTAAACCGTCCGTGCCAGTCCACTTCGATGCTTCAGTGAAGGACG  
AACTGTAA

ATGAAATACCTATTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCAACC  
AGCGATGGCCCAGCTGCAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAGAGCC  
TGTCATCACATGCACTGTCTCAGGGTTCTCATTAACCAGTTATGGTGTAAGCTGG  
GTTCCGCCAGCCTCCAAGAAAGGGTCTGGAGTGGCTGGGAGTAATATGGGAAGACGG  
GAGCACAATTATCATTACGTCTCATATCCAGACTGAGCATCAACAAGGATAACT  
CCAAGAGCCAAGTTTTCTTAAACTGAACAGTCTGCAAACCTGATGACACAGCCACG  
TACTACTGTGCCAAACCCCACTACGGTAGCAGCAACGTGGGGGCTATGGAATACTG  
GGGTCAAGGAACCTCGGTACCCGTCTCCTCAGGTGGAGGCGGTTTCAGGCGGAGGTG  
GCTCTGGCGGTGGCGGATCGGACATCGAGCTACCCAGTCTCCAGCCTCCCTAACT  
GCATCTGTGGGAGAACTGTCACCATCACCTGTGAGCAAGTGAAAATATTTACAG  
TTATGTAGCATGGTATCAGCAGAAACAGGGAAAATCTCCTCAGTTCCTGGTCTATA  
ATGCAAAATCCTTAGCAGAGGGTGTGCCATCAAGGTTCAAGTGGCAGTGGATCAGGC  
ACACAGTTTTCTCTGAAGATCAACAGCCTGCAGCCTGAAGATTTTGGGAATTATTA  
CTGTCAACATCATTATGTTAGTCCGTGGACGTTCCGTGGAGGCACCAAGCTCGAGA  
TCAAGGCACCTGCTGCCTCCCCGGCAGACGCTAAGGAATCTGCAGCTGCCAAGTTC  
GAGCGGCAGCACATGGACTCTGGCAACTCCCCCAGCAGCAGTCCAACTACTGCAA  
CCTGATGATGTGCTGCCGGAAGATGACCCAGGGGAAATGCAAGCCAGTGAACACCT  
TTGTGCATGAGTCCCTGGCCGATGTTAAGGCCGTGTGCTCCCAGAAGAAAGTCACT  
TGCAAGAATGGGCAGACCAACTGCTACCAGAGCAAATCCACCATGCGCATCACAGA  
CTGCCGCGAGACTGGCAGTCCAAGTACCCCAACTGCGCCTACAAGACCACCCAGG  
TGGAGAAACACATCATAGTGGCTTGTGGCGGTAAACCGTCCGTGCCAGTCCACTTC  
GATGCTTCAGTGAAGGACGAACTGTAA

ATGAAATACCTATTGCCTACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCAACC  
AGCGATGGCCCAGCTGCAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAGAGCC  
TGTCATCACATGCACTGTCTCAGGGTTCTCATTAACCAGTTATGGTGTAAGCTGG  
GTTCCGCCAGCCTCCAAGAAAGGGTCTGGAGTGGCTGGGAGTAATATGGGAAGACGG  
GAGCACAATTATCATTACAGTCTCATATCCAGACTGAGCATCAACAAGGATAACT  
CCAAGAGCCAAGTTTTCTTAAACTGAACAGTCTGCAAAGTATGACACAGCCACG  
TACTACTGTGCCAAACCCCACTACGGTAGCAGCAACGTGGGGGCTATGGAATACTG  
GGGTCAAGGAACCTCGGTACCGTCTCCTCAGGTGGAGGCGGTTTCAGGCGGAGGTG  
GCTCTGGCGGTGGCGGATCGGACATCGAGCTCAGGAGTCTCCAGCCTCCCTAACT  
GCATCTGTGGGAGAACTGTCAACATCACCTGTGAGCAAGTGAATAATTTACAG  
TTATGTAGCATGGTATCAGCAGAAACAGGGAAAATCTCCTCAGTTCCTGGTCTATA  
ATGCAAAATCCTTAGCAGAGGGTGTGCCATCAAGGTTTCAGTGGCAGTGGATCAGGC  
ACACAGTTTTCTCTGAAGATCAACAGCCTGCAGCCTGAAGATTTTGGGAATTATTA  
CTGTCAACATCATTATGTTAGTCCGTGGACGTTTCGGTGGAGGCACCAAGCTCGAGA  
TCAAGCGCAAGGAATCTGCAGCTGCCAAGTTTCGAGCGGCAGCACATGGACTCTGGC  
AACTCCCCCAGCAGCAGCTCCAACTACTGCAACCTGATGATGTGCTGCCGGAAGAT  
GACCCAGGGGAAATGCAAGCCAGTGAACACCTTTGTGCATGAGTCCCTGGCCGATG  
TTAAGGCCGTGTGCTCCCAGAAGAAAGTCACTTGCAAGAATGGGCAGACCAACTGC  
TACCAGAGCAAATCCACCATGCGCATCACAGACTGCCGCGAGACTGGCAGCTCCAA  
GTACCCCAACTGCGCCTACAAGACCAACCAGGTGGAGAAACACATCATAGTGGCTT  
GTGGCGGTAAACCGTCCGTGCCAGTCCACTTCGATGCTTCAGTGAAGGACGAACTG  
TAA

FIGURE 33




Fig. 34



## COMPOUNDS FOR TARGETING

The present invention relates to compounds, some of which may be directly or indirectly cytotoxic combinations of compounds, that have a high avidity for, and can be targeted to, selected cells.

### Background and Prior Art

The cell-specific targeting of compounds which are directly, or indirectly, cytotoxic has been proposed as a way to combat diseases such as cancer. Bagshawe and his co-workers have disclosed (Bagshawe (1987) *Br. J. Cancer* 56, 531; Bagshawe *et al* (1988) *Br. J. Cancer* 58, 700; WO 88/07378) conjugated compounds comprising an antibody or part thereof and an enzyme, the antibody being specific to tumour cell antigens and the enzyme acting to convert an innocuous pro-drug into a cytotoxic compound. The cytotoxic compounds were alkylating agents, eg a benzoic acid mustard released from para-N-bis(2-chloroethyl)aminobenzoyl glutamic acid by the action of *Pseudomonas sp.* CPG2 enzyme.

An alternative system using different pro-drugs has been disclosed (WO 91/11201) by Epenetos and co-workers. The cytotoxic compounds were cyanogenic monosaccharides or disaccharides, such as the plant compound amygdalin, which release cyanide upon the action of a  $\beta$ -glucosidase and hydroxynitrile lyase.

In a further alternative system, the use of antibody-enzyme conjugates containing the enzyme alkaline phosphatase in conjunction with the pro-drug etoposide 4'-phosphate or 7-(2'-aminoethyl phosphate)mitomycin or a combination thereof have been disclosed (EP 0 302 473; Senter *et al* (1988) *Proc. Natl. Acad. Sci. USA* 85, 4842).

Rybak and co-workers have disclosed (Rybak *et al* (1991) *J. Biol. Chem.* 266, 21202; WO 91/16069) the cytotoxic potential of a monomeric pancreatic ribonuclease when injected directly into *Xenopus* oocytes and the cytotoxic potential of monomeric RNase coupled to human transferrin or antibodies directed against the transferrin receptor. The monomeric RNase hybrid proteins were cytotoxic to human erythroleukaemia cells *in vitro*.

Other approaches are the *in vivo* application of streptavidin conjugated antibodies followed, after an appropriate period, by radioactive biotin (Hnatowich *et al* (1988) *J. Nucl. Med.* 29, 1428-1434), or injection of a biotinylated mAb followed by radioactive streptavidin (Paganelli *et al* (1990) *Int. J. Cancer* 45, 1184-1189). A pilot radioimmunolocalisation study in non-small cell lung carcinomas was conducted with encouraging results (Kalofonos *et al* (1990) *J. Nucl. Med.* 31, 1791-1796).

Apart from these examples, it is rather more common to see biotinylated antibodies and streptavidin-enzyme conjugates which are used in enzyme-linked immunosorbent assays.

These previous systems have used relatively large antibody-enzyme or antibody-streptavidin or antibody-biotin conjugates and may comprise portions of non-mammalian origin which are highly immunoreactive.

Rapid penetrance (Yokota *et al* (1992) *Cancer Res.* 52, 3402-3408) and rapid clearance (Colcher *et al* (1990) *J. Natl. Cancer Inst.* 82, 1191-1197) has been demonstrated for single chain Fv antibody fragments (ScFv).

In using the cell-specific reagents aforementioned in a therapeutically useful situation one of the requirements that needs to be met is for the

cell-specific reagent to accumulate to a sufficiently higher level at the target cell than at other cells. A further requirement is that a directly or indirectly cytotoxic reagent is carried to the target cell, and it is preferred that the said cytotoxic reagent is of high potency.

5

We have now devised improved systems at least some of which exhibit higher avidities to the selected target cells, and make use of novel, potent directly or indirectly cytotoxic agents.

## 10 Summary of Invention

A first aspect of the invention provides a compound comprising a target cell-specific portion and a cytotoxic portion characterised in that the cytotoxic portion has nucleolytic activity.

15

Suitably, as disclosed below, the cytotoxic portion may have ribonucleolytic activity or it may have DNA endonucleolytic activity.

20

One aspect of the present invention provides a compound comprising a target cell-specific portion and a directly or indirectly cytotoxic portion, characterised in that the target cell-specific portion recognises the target cell with high avidity.

25

A further aspect of the present invention provides a compound comprising a target cell-specific portion and a directly or indirectly cytotoxic portion characterised in that the cytotoxic portion is a sub-unit of an oligomer provided that, if the sub-unit is complexed with another sub-unit of the said oligomer then the said other sub-unit is the cytotoxic portion of a second compound of the invention.

30

A further aspect of the present invention provides a compound of at least two molecules each comprising a target cell-specific portion and a further portion wherein the molecules are complexed to one another via their further portions.

5

A further aspect of the present invention provides a compound comprising an oligomeric complex of at least two molecules each comprising a target cell-specific portion wherein the molecules are complexed to one another via their cytotoxic portions.

10

A further aspect of the present invention provides a compound comprising a target cell-specific portion and a directly or indirectly cytotoxic portion characterised in that the cytotoxic portion contains a binding site for a small-molecule wherein the said small-molecule binding site binds but does not modify catalytically the said small molecule.

15

A further aspect of the present invention provides a compound comprising a target cell-specific portion and a directly or indirectly cytotoxic portion characterised in that the target cell-specific portion comprises two or more binding sites for the target cell, wherein the target cell-specific portion is not an antibody, or bivalent fragment thereof, having respective arms which recognise the same entity as one another.

20

A further aspect of the present invention provides a compound comprising a target cell-specific portion and a cytotoxic portion characterised in that the cytotoxic portion has DNA endonucleolytic activity.

25

A further aspect of the invention provides a compound comprising a mediator portion and a directly or indirectly cytotoxic portion.

30

By "mediator portion" we mean the portion of the compound that recognises a target cell-specific molecule. The target cell-specific molecule may be a further compound of any of the appropriate preceding aspects of the present invention or it may be a target cell-specific molecule known in the art or it may be a derivative thereof capable of recognition by the mediator portion.

By "high avidity" we mean that the target cell-specific portion recognises the target cell with a binding constant of at least  $K_d = 10^{-9}M$ , suitably  $K_d = 10^{-10}M$ , more suitably  $K_d = 10^{-11}M$ , more suitably still  $K_d = 10^{-12}M$ , preferably  $K_d = 10^{-15}M$ , more preferably  $K_d = 10^{-18}M$ , more preferably still  $K_d = 10^{-21}M$ , yet even more preferably  $K_d = 10^{-24}M$ , and in further preference  $K_d = 10^{-27}M$  or even  $K_d = 10^{-30}M$ .

By "target cell specific" portion we mean the portion of the compound which comprises one or more binding sites which recognise and bind to entities on the target cell. The said entities are expressed predominantly, and preferably exclusively, on the said target cell. The target cell specific portion may contain one or more binding sites for different entities expressed on the same target cell type, or one or more binding sites for different entities expressed on two or more different target cell types.

By a "directly cytotoxic agent" we mean an agent which in itself is toxic to the cell if it is to reach, and preferably enter the said cell.

By an "indirectly cytotoxic agent" we mean an agent which in itself may or may not be non-toxic, but which can bind specifically to a cytotoxic compound, or can bind specifically to a compound which can be converted into a cytotoxic compound by the action of a further reagent.

- The entity which is recognised may be any suitable entity which is expressed by tumour cells, virally-infected cells, pathogenic microorganisms, cells introduced as part of gene therapy or even normal cells of the body which, for whatever reason, one wishes to target, but which is not expressed, or at least not with such frequency, in cells which one does not wish to target. The entity which is recognised will often be an antigen. Examples of antigens include those listed in Table 1 below. Monoclonal antibodies which will bind to many of these antigens are already known (for example those given in the Table) but in any case, with today's techniques in relation to monoclonal antibody technology, antibodies can be prepared to most antigens. The antigen-specific portion may be a part of an antibody (for example a Fab fragment) or a synthetic antibody fragment (for example a single chain Fv fragment [ScFv]). Suitable monoclonal antibodies to selected antigens may be prepared by known techniques, for example those disclosed in "*Monoclonal Antibodies: A manual of techniques*", H Zola (CRC Press, 1988) and in "*Monoclonal Hybridoma Antibodies: Techniques and Applications*", J G R Hurrell (CRC Press, 1982).
- 20 The variable heavy ( $V_H$ ) and variable light ( $V_L$ ) domains of the antibody are involved in antigen recognition, a fact first recognised by early protease digestion experiments. Further confirmation was found by "humanisation" of rodent antibodies. Variable domains of rodent origin may be fused to constant domains of human origin such that the resultant antibody retains the antigenic specificity of the rodent parented antibody (Morrison *et al* (1984) *Proc. Natl. Acad. Sci. USA* 81, 6851-6855).

That antigenic specificity is conferred by variable domains and is independent of the constant domains is known from experiments involving the bacterial expression of antibody fragments, all containing one or more

variable domains. These molecules include Fab-like molecules (Better *et al* (1988) *Science* **240**, 1041); Fv molecules (Skerra *et al* (1988) *Science* **240**, 1038); single-chain Fv (ScFv) molecules where the V<sub>H</sub> and V<sub>L</sub> partner domains are linked via a flexible oligopeptide (Bird *et al* (1988) *Science* **242**, 423; Huston *et al* (1988) *Proc. Natl. Acad. Sci. USA* **85**, 5879) and single domain antibodies (dAbs) comprising isolated V domains (Ward *et al* (1989) *Nature* **341**, 544). A general review of the techniques involved in the synthesis of antibody fragments which retain their specific binding sites is to be found in Winter & Milstein (1991) *Nature* **349**, 293-299.

By "ScFv molecules" we mean molecules wherein the V<sub>H</sub> and V<sub>L</sub> partner domains are linked via a flexible oligopeptide.

15 Chimaeric antibodies are discussed by Neuberger *et al* (1988, *8th International Biotechnology Symposium Part 2*, 792-799).

Suitably prepared non-human antibodies can be "humanized" in known ways, for example by inserting the CDR regions of mouse antibodies into the framework of human antibodies.

The advantages of using antibody fragments, rather than whole antibodies, are several-fold. The smaller size of the fragments allows for rapid clearance, and may lead to improved tumour to non-tumour ratios. Fab, Fv, ScFv and dAb antibody fragments can all be expressed in and secreted from *E. coli*, thus allowing the facile production of large amounts of the said fragments.

Whole antibodies, and F(ab')<sub>2</sub> fragments are "bivalent". By "bivalent" we mean that the said antibodies and F(ab')<sub>2</sub> fragments have two antigen

combining sites. In contrast, Fab, Fv, ScFv and dAb fragments are monovalent, having only one antigen combining site.

Alternatively, the entity which is recognised may or may not be antigenic but can be recognised and selectively bound to in some other way. For example, it may be a characteristic cell surface receptor such as the receptor for melanocyte-stimulating hormone (MSH) which is expressed in high number in melanoma cells. The cell-specific portion may then be a compound or part thereof which specifically binds to the entity in a non-immune sense, for example as a substrate or analogue thereof for a cell-surface enzyme or as a messenger.

Preferably, the high avidity target cell specific portion comprises two or more different binding sites for the target cell.

15

The different binding sites for the target cell may or may not be two or more different antibodies, or fragments thereof, which are directed to different entities expressed on the target cell. Alternatively, the different binding sites for the target cell may recognise and selectively bind the cell in some other, non-immune sense.

20

A further alternative is that one or more of the binding sites is an antibody, or part thereof, and that one or more of the binding sites for the target cell recognise and selectively bind the cell in some other, non-immune sense.

25

A compound which has binding sites for two or more target cell-specific entities may be more specific for binding to the said target cell, and a compound which has more than one of each of the different binding sites may bind to the said target cell with greater avidity. In combining two or

30



more binding sites, which in themselves may be of high specificity but low affinity, it will be possible to generate in the compound of the invention a higher affinity for the target cell whilst retaining the specificity of the binding sites.

Table 11. Tumour Associated Antigens

5	<u>Antigen</u>	<u>Antibody</u>	<u>Existing Uses</u>
	Carcino-embryonic Antigen	{C46 (Amersham) {85A12 (Unipath)	Imaging & Therapy of colon/rectum tumours.
	Placental Alkaline Phosphatase	H17E2 (ICRF, Travers & Bodmer)	Imaging & Therapy of testicular and ovarian cancers.
10	Pan Carcinoma	NR-LU-10 (NeoRx Corporation)	Imaging & Therapy of various carcinomas incl. small cell lung cancer.
	Polymorphic Epithelial Mucin (Human milk fat globule	HMFG1 (Taylor-Papadimitriou, ICRF)	Imaging & Therapy of ovarian cancer, pleural effusions.
15	Human milk mucin core protein	SM-3(IgG1) <sup>1</sup>	Diagnosis, Imaging & Therapy of breast cancer

- |   |  |                          |   |
|---|--|--------------------------|---|
|   | $\beta$ -human Chorionic Gonadotropin              | W14                      | Targeting of enzyme (CPG2) to human xenograft choriocarcinoma in nude mice. (Searle <i>et al</i> (1981) <i>Br. J. Cancer</i> <b>44</b> , 137-144) |
|   | A Carbohydrate on Human Carcinomas                 | L6 (IgG2a) <sup>2</sup>  | Targeting of alkaline phosphatase. (Senter <i>et al</i> (1988) <i>Proc. Natl. Acad. Sci. USA</i> <b>85</b> , 4842-4846)                           |
| 5 | CD20 Antigen on B Lymphoma (normal and neoplastic) | 1F5 (IgG2a) <sup>3</sup> | Targeting of alkaline phosphatase. (Senter <i>et al</i> (1988) <i>Proc. Natl. Acad. Sci. USA</i> <b>85</b> , 4842-4846)                           |
- <sup>1</sup>Burchell *et al* (1987) *Cancer Res.* **47**, 5476-5482
- <sup>2</sup>Hellström *et al* (1986) *Cancer Res.* **46**, 3917-3923
- 10 <sup>3</sup>Clarke *et al* (1985) *Proc. Natl. Acad. Sci. USA* **82**, 1766-1770

Other antigens include alphafoetoprotein, Ca-125 and prostate specific antigen.

## 2. Immune Cell Antigens

5	Pan T Lymphocyte Surface Antigen (CD3)	OKT-3 (Ortho)	As anti-rejection therapy for kidney transplants.
	B-lymphocyte Surface Antigen (CD22)	RFB4 (Janossy, Royal Free Hospital)	Immunotoxin therapy of B cell lymphoma.
10	Pan T lymphocyte Surface Antigen (CD5)	H65 (Bodmer, Knowles ICRF, Licensed to Xoma Corp., USA)	Immunotoxin treatment of Acute Graft versus Host disease, Rheumatoid Arthritis.

## 3. Infectious Agent-Related Antigens

	Mumps virus-related	Anti-mumps polyclonal antibody	Antibody conjugated to Diphtheria toxin for treatment of mumps.
15	Hepatitis B Surface Antigen	Anti HBs Ag	Immunotoxin against Hepatoma.

It is preferable that the two portions of the compound of the invention are produced as a fusion compound by recombinant DNA techniques whereby a length of DNA comprises respective regions encoding the two portions of the compound of the invention either adjacent one another or separated by a region encoding a linker peptide which does not destroy the desired

- properties of the compound. The benefits in making the compound of the invention using recombinant DNA techniques are several fold. Firstly, it enables a high degree of precision with which the two portions of the compound can be joined together. Secondly, the construction of compounds which are "hetero-oligomeric" can be controlled by the expression of the different recombinant DNA molecules encoding each of the different type of subunit of the "hetero-oligomer" in the same host cell.
- 10 By "hetero-oligomer" we mean those compounds in which two or more different cell-specific portions are joined to either the same or to different subunits which are capable of oligomerisation. The expression, in the same host cell of two compounds, A and B, each with different target cell specific portions but with a common second portion capable of oligomerisation will result in a mixed population of compounds. For example, if the common second portion is capable of dimerisation, three potential compounds will be produced:  $A_2$ , AB and  $B_2$ , in a ratio of 1:2:1, respectively.
- 20 The separation of the desired compound with each of the different cell specific portions, that is AB, can be achieved by two step affinity chromatography.
- 25 Application of the mixture of compounds to an affinity column specific for A will result in the binding of  $A_2$  and AB. These compounds are eluted from this first column, and then applied to an affinity column specific for B. This will result in AB, but not  $A_2$ , being bound to the column. Finally, the desired product AB, can be eluted.
- 30 Of course, the order in which the affinity columns are used is not

important.

The same principle of separating those compounds with two or more different binding sites can be applied to the purification of the desired  
5 compounds from mixtures of other hetero-oligomers.

Conceivably, the two portions of the compound may overlap wholly or partly.

10 The DNA is then expressed in a suitable host to produce a polypeptide comprising the compound of the invention. Thus, the DNA encoding the polypeptide constituting the compound of the invention may be used in accordance with known techniques, appropriately modified in view of the teachings contained herein, to construct an expression vector, which is  
15 then used to transform an appropriate host cell for the expression and production of the polypeptide of the invention. Such techniques include those disclosed in US Patent Nos. 4,440,859 issued 3 April 1984 to Rutter *et al*, 4,530,901 issued 23 July 1985 to Weissman, 4,582,800 issued 15 April 1986 to Crawl, 4,677,063 issued 30 June 1987 to Mark *et al*,  
20 4,678,751 issued 7 July 1987 to Goeddel, 4,704,362 issued 3 November 1987 to Itakura *et al*, 4,710,463 issued 1 December 1987 to Murray, 4,757,006 issued 12 July 1988 to Toole, Jr. *et al*, 4,766,075 issued 23 August 1988 to Goeddel *et al* and 4,810,648 issued 7 March 1989 to Stalker, all of which are incorporated herein by reference.

25

The DNA encoding the polypeptide constituting the compound of the invention may be joined to a wide variety of other DNA sequences for introduction into an appropriate host. The companion DNA will depend upon the nature of the host, the manner of the introduction of the DNA  
30 into the host, and whether episomal maintenance or integration is desired.

Generally, the DNA is inserted into an expression vector, such as a plasmid, in proper orientation and correct reading frame for expression.

If necessary, the DNA may be linked to the appropriate transcriptional and translational regulatory control nucleotide sequences recognised by the

5 desired host, although such controls are generally available in the expression vector. The vector is then introduced into the host through standard techniques. Generally, not all of the hosts will be transformed by the vector. Therefore, it will be necessary to select for transformed host cells. One selection technique involves incorporating into the  
10 expression vector a DNA sequence, with any necessary control elements, that codes for a selectable trait in the transformed cell, such as antibiotic resistance. Alternatively, the gene for such selectable trait can be on another vector, which is used to co-transform the desired host cell.

15 Host cells that have been transformed by the recombinant DNA of the invention are then cultured for a sufficient time and under appropriate conditions known to those skilled in the art in view of the teachings disclosed herein to permit the expression of the polypeptide, which can then be recovered.

20

Many expression systems are known, including bacteria (for example *E. coli* and *Bacillus subtilis*), yeasts (for example *Saccharomyces cerevisiae*), filamentous fungi (for example *Aspergillus*), plant cells, animal cells and insect cells.

25

Those vectors that include a replicon such as a procaryotic replicon can also include an appropriate promoter such as a procaryotic promoter capable of directing the expression (transcription and translation) of the genes in a bacterial host cell, such as *E. coli*, transformed therewith.

30

A promoter is an expression control element formed by a DNA sequence that permits binding of RNA polymerase and transcription to occur. Promoter sequences compatible with exemplary bacterial hosts are typically provided in plasmid vectors containing convenient restriction sites  
5 for insertion of a DNA segment of the present invention.

Typical procaryotic vector plasmids are pUC18, pUC19, pBR322 and pBR329 available from Biorad Laboratories, (Richmond, CA, USA) and pTrc99A and pKK223-3 available from Pharmacia, Piscataway, NJ, USA.  
10

A typical mammalian cell vector plasmid is pSVL available from Pharmacia, Piscataway, NJ, USA. This vector uses the SV40 late promoter to drive expression of cloned genes, the highest level of expression being found in T antigen-producing cells, such as COS-1 cells.  
15

An example of an inducible mammalian expression vector is pMSG, also available from Pharmacia. This vector uses the glucocorticoid-inducible promoter of the mouse mammary tumour virus long terminal repeat to drive expression of the cloned gene.  
20

Useful yeast plasmid vectors are pRS403-406 and pRS413-416 and are generally available from Stratagene Cloning Systems, La Jolla, CA 92037, USA. Plasmids pRS403, pRS404, pRS405 and pRS406 are Yeast Integrating plasmids (YIps) and incorporate the yeast selectable markers  
25 *his3*, *trp1*, *leu2* and *ura3*. Plasmids pRS413-416 are Yeast Centromere plasmids (YCps).

A variety of methods have been developed to operatively link DNA to vectors via complementary cohesive termini. For instance,  
30 complementary homopolymer tracts can be added to the DNA segment to



be inserted to the vector DNA. The vector and DNA segment are then joined by hydrogen bonding between the complementary homopolymeric tails to form recombinant DNA molecules.

- 5 Synthetic linkers containing one or more restriction sites provide an alternative method of joining the DNA segment to vectors. The DNA segment, generated by endonuclease restriction digestion as described earlier, is treated with bacteriophage T4 DNA polymerase or *E. coli* DNA polymerase I, enzymes that remove protruding, 3'-single-stranded termini  
10 with their 3'-5'-exonucleolytic activities, and fill in recessed 3'-ends with their polymerizing activities.

The combination of these activities therefore generates blunt-ended DNA segments. The blunt-ended segments are then incubated with a large  
15 molar excess of linker molecules in the presence of an enzyme that is able to catalyze the ligation of blunt-ended DNA molecules, such as bacteriophage T4 DNA ligase. Thus, the products of the reaction are DNA segments carrying polymeric linker sequences at their ends. These DNA segments are then cleaved with the appropriate restriction enzyme  
20 and ligated to an expression vector that has been cleaved with an enzyme that produces termini compatible with those of the DNA segment.

Synthetic linkers containing a variety of restriction endonuclease sites are commercially available from a number of sources including International  
25 Biotechnologies Inc, New Haven, CN, USA.

A desirable way to modify the DNA encoding the polypeptide of the invention is to use the polymerase chain reaction as disclosed by Saiki *et al* (1988) *Science* 239, 487-491.

In this method the DNA to be enzymatically amplified is flanked by two specific oligonucleotide primers which themselves become incorporated into the amplified DNA. The said specific primers may contain restriction endonuclease recognition sites which can be used for cloning into  
5 expression vectors using methods known in the art.

Exemplary genera of yeast contemplated to be useful in the practice of the present invention are *Pichia*, *Saccharomyces*, *Kluyveromyces*, *Candida*, *Torulopsis*, *Hansenula*, *Schizosaccharomyces*, *Citeromyces*, *Pachysolen*,  
10 *Debaromyces*, *Metschnikowia*, *Rhodospiridium*, *Leucosporidium*, *Botryosascus*, *Sporidiobolus*, *Endomycopsis*, and the like. Preferred genera are those selected from the group consisting of *Pichia*, *Saccharomyces*, *Kluyveromyces*, *Yarrowia* and *Hansenula*. Examples of *Saccharomyces* are *Saccharomyces cerevisiae*, *Saccharomyces italicus* and *Saccharomyces*  
15 *rouxii*. Examples of *Kluyveromyces* are *Kluyveromyces fragilis* and *Kluyveromyces lactis*. Examples of *Hansenula* are *Hansenula polymorpha*, *Hansenula anomala* and *Hansenula capsulata*. *Yarrowia lipolytica* is an example of a suitable *Yarrowia* species.

20 Methods for the transformation of *S. cerevisiae* are taught generally in EP 251 744, EP 258 067 and WO 90/01063, all of which are incorporated herein by reference.

Suitable promoters for *S. cerevisiae* include those associated with the  
25 *PGK1* gene, *GAL1* or *GAL10* genes, *CYC1*, *PHO5*, *TRP1*, *ADH1*, *ADH2*, the genes for glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, triose phosphate isomerase, phosphoglucose isomerase, glucokinase,  $\alpha$ -mating factor pheromone, a-mating factor pheromone, the *PRB1* promoter, the *GUT2* promoter, and  
30 hybrid promoters involving hybrids of parts of 5' regulatory regions with

parts of 5' regulatory regions of other promoters or with upstream activation sites (eg the promoter of EP-A-258 067).

5 The transcription termination signal is preferably the 3' flanking sequence of a eukaryotic gene which contains proper signals for transcription termination and polyadenylation. Suitable 3' flanking sequences may, for example, be those of the gene naturally linked to the expression control sequence used, ie may correspond to the promoter. Alternatively, they may be different in which case the termination signal of the *S. cerevisiae*  
10 *AHD1* gene is preferred.

The present invention also relates to a host cell transformed with a polynucleotide vector construct of the present invention. The host cell can be either procaryotic or eucaryotic. Bacterial cells are preferred  
15 procaryotic host cells and typically are a strain of *E. coli* such as, for example, the *E. coli* strains DH5 available from Bethesda Research Laboratories Inc., Bethesda, MD, USA, and RR1 available from the American Type Culture Collection (ATCC) of Rockville, MD, USA (No ATCC 31343). Preferred eucaryotic host cells include yeast and  
20 mammalian cells, preferably vertebrate cells such as those from a mouse, rat, monkey or human fibroblastic cell line. Preferred eucaryotic host cells include Chinese hamster ovary (CHO) cells available from the ATCC as CCL61, NIH Swiss mouse embryo cells NIH/3T3 available from the ATCC as CRL 1658 and monkey kidney-derived COS-1 cells available  
25 from the ATCC as CRL 1650.

Transformation of appropriate cell hosts with a DNA construct of the present invention is accomplished by well known methods that typically depend on the type of vector used. With regard to transformation of  
30 procaryotic host cells, see, for example, Cohen *et al*, *Proc. Natl. Acad.*

Sci. USA, 69: 2110 (1972); and Sambrook *et al*, *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1989). Transformation of yeast cells is described in Sherman *et al*, *Methods In Yeast Genetics, A Laboratory Manual*, Cold Spring Harbor, NY (1986). The method of Beggs, *Nature*, 275: 104-109 (1978) is also useful. With regard to vertebrate cells, reagents useful in transfecting such cells, for example calcium phosphate and DEAE-dextran or liposome formulations, are available from Stratagene Cloning Systems, or Life Technologies Inc, Gaithersburg, MD 20877, USA.

10

Successfully transformed cells, ie cells that contain a DNA construct of the present invention, can be identified by well known techniques. For example, cells resulting from the introduction of an expression construct of the present invention can be grown to produce the polypeptide of the invention. Cells can be harvested and lysed and their DNA content examined for the presence of the DNA using a method such as that described by Southern, *J. Mol. Biol.*, 98: 503 (1975) or Berent *et al*, *Biotech.*, 3: 208 (1985). Alternatively, the presence of the protein in the supernatant can be detected using antibodies as described below.

20

In addition to directly assaying for the presence of recombinant DNA, successful transformation can be confirmed by well known immunological methods when the recombinant DNA is capable of directing the expression of the protein. For example, cells successfully transformed with an expression vector produce proteins displaying appropriate antigenicity. Samples of cells suspected of being transformed are harvested and assayed for the protein using suitable antibodies.

25

Thus, in addition to the transformed host cells themselves, the present invention also contemplates a culture of those cells, preferably a

30

monoclonal (clonally homogeneous) culture, or a culture derived from a culture, in a nutrient medium. Preferably, the culture also contains the protein.

- 5 Nutrient media useful for culturing transformed host cells are well known in the art and can be obtained from several commercial sources.

Alternatively, the target-cell specific and second portions of the compound of the invention are linked together by any of the conventional ways of cross-linking polypeptides, such as those generally described in O'Sullivan  
10 *et al Anal. Biochem.* (1979) 100, 100-108. For example, the antibody portion may be enriched with thiol groups and the enzyme portion reacted with a bifunctional agent capable of reacting with those thiol groups, for example the N-hydroxysuccinimide ester of iodoacetic acid (NHIA) or N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP). Amide and thioether  
15 bonds, for example achieved with m-maleimidobenzoyl-N-hydroxysuccinimide ester, are generally more stable *in vivo* than disulphide bonds.

- 20 Some of the various compounds of the invention are illustrated diagrammatically in Figure 1. C and D are the target cell-specific portions, and X is the cytotoxic portion. Of course, X may form higher order oligomers than those illustrated for example trimers, tetramers, pentamers, hexamers.

25

In Figure 1(a) to 1(d) C and D are shown binding to entities on either the same, or different cells.

- In one embodiment of the invention, C and D recognise different  
30 molecules on the same target cell wherein the molecules on the same

- target cell are not confined to that cell type but may occur on a few other cell types. In particular, C may recognise molecules on cell types I, II and III, whereas D may recognise molecules on cell types I, IV and V. Thus a compound of the invention comprising C and D as the target cell-specific portion will have greater specificity for cell type I compared with cell types II, III and IV. This aspect of the invention is particularly helpful, as there have been very few completely target cell-specific molecules discovered, whereas molecules which occur on a few cell types, and which are useful in this aspect of the invention, are well known.
- Such molecules are usually cell-surface antigens for which cross-reactive antibodies are known. Examples of such molecules are given in Table 2.

**Table 2**

<u>Antigen</u>	<u>Cell-type</u>	<u>Antibody</u>
CD9	Pre-B cells, monocytes, platelets	MM2/57 (IgG2b, mouse)
CALLA	Lymphoid progenitor cells, granulocytes	B-E3 (IgG2a, mouse)
CD13	Myeloid monocytes, granulocytes	B-F10 (IgG1, mouse)
CD24	B-cells, granulocytes	ALB-9 (IgG1, mouse)
CD61	Platelets, megakaryocytes	PM 6/13 (IgG1, mouse)

The antibodies described in Table 2 are generally available from Serotec, Oxford, OX5 1BR, UK.

Preferably, the cytotoxic portion of the compound of the invention is capable of oligomerisation. Attachment of the target-cell specific portion to a cytotoxic portion capable of oligomerisation provides a method for increasing the number of binding sites to the target cell. For example, if  
5 the target cell-specific portion is joined to a portion capable of forming a dimer then the number of target cell-specific binding sites is two; if the target cell-specific portion is joined to a portion capable of forming a tetramer then the number of target cell-specific binding sites is four. The number of target cell-specific binding sites is greater than one and the  
10 compounds may therefore have a greater avidity for the target cell than do compounds which only have one target cell-specific binding site.

It is preferable for the cytotoxic portion of the compound of the invention capable of oligomerisation to contain no interchain disulphide bonds nor  
15 intrachain disulphide bonds; to be well characterised; to be non-toxic; to be stable; to be amenable to preparation in a form suitable for pre-clinical or clinical use or be in pre-clinical or clinical use; and for the subunit monomers to have a high affinity for each other, that is they contain one or more subunit binding sites.

20

Preferably, each subunit of the cytotoxic portion of the compound of the invention contains a binding site for a small molecule, the small molecule being capable of being conjugated to any from the following compounds: radioactive compound; spin-labelled compound; drug; pro-drug;  
25 radionuclide; protein including enzyme; antibody; or toxin.

In a preferred embodiment of the invention, the cytotoxic portion is streptavidin. Streptavidin is a homotetrameric molecule of  $M_r = 60000$  (subunit  $M_r = 15000$ ) and is produced by *Streptomyces*. Streptavidin  
30 binds four molecules of the water-soluble vitamin biotin with high

specificity and affinity ( $K_d = 10^{-15}M$ ) although isolated subunits possess a very much lower affinity for biotin ( $K_d = 10^{-8}M$ ). Each subunit of streptavidin has a tightly-packed "core", with relatively unstructured amino- and carboxyl-terminal extensions. These extensions are believed to contribute to the formation of higher order aggregates of streptavidin. Many commercial forms of streptavidin are extensively proteolysed, have lost their unstructured extensions, and form stable tetramers (Bayer *et al* (1989) *Biochem J.* **259**, 369-376; Bayer *et al* (1990) *Methods Enzymol.* **184**, 51-67). The mature form of the protein has been the subject of recent research and is becoming increasingly well characterised (Gitlin *et al* (1988) *Biochem J.* **256**, 279-282; Gitlin *et al* (1990) *Biochem J.* **269**, 527-530; Sano & Cantor (1990) *J. Biol. Chem.* **265**, 3369-3373) and the gene has been cloned and sequenced (Argarana *et al* (1986) *Nucl. Acids Res.* **14**, 1871-1872) and expressed in *E. coli* (Sano & Cantor (1990) *Proc. Natl. Acad. Sci. USA* **87**, 142-146). A modified form of the gene is available commercially from British Bio-technology Ltd, Oxford, UK.

Of course, for the invention to work the cytotoxic portion may comprise intact streptavidin, or it may comprise a fragment or fragments of streptavidin retaining at least the biotin- and subunit-binding sites.

Of course, the cytotoxic portion may comprise other molecules which bind biotin with high affinity, such as intact avidin, or it may comprise a fragment or fragments of avidin retaining at least the biotin- and subunit-binding sites. A comparison of avidin and streptavidin is made in Table 3. As avidin is naturally glycosylated, then it may be desirable to express the DNA encoding the compound of the invention in a eukaryotic cell such as yeast, mammalian or insect cell.



	<u>Avidin</u>	<u>Streptavidin</u>
Source	Tissues and egg-whites of birds, reptiles and amphibia	<i>Streptomyces avidinii</i>
5 Glycoprotein	yes	no
pI	10	5
M <sub>r</sub> (subunit)	67,000	60,000
Oligomeric state	Tetramer	Tetramer

- 10 By "subunit-binding sites" we mean those parts of the monomers that are necessary for the monomers to combine with one or more other monomers to produce an oligomer.

15 Biotin has an extremely high affinity for streptavidin ( $K_d = 10^{-15}M$ ) and at the same time is small enough to diffuse rapidly through most tissues in the body. Some of the biotin conjugates useful in the invention are known in the art, and it is preferred that the biotin is conjugated via a flexible linker arm to reduce any steric hindrance to the binding of the biotin portion of the conjugate to streptavidin or avidin.

20

Examples of biotin conjugates useful in the invention are biotinylated growth factors and cytokines such as TNF $\alpha$ -biotin and EGF-biotin which are generally available from Boehringer Mannheim, Mannheim, Germany, and biotin-alkaline phosphatase, biotin-fluorescein, biotin-peroxidase and  
25 other conjugates generally available from Calbiochem-Novabiochem, Nottingham, UK. Activated biotin reagents, suitable for conjugating to other molecules, are generally available from Fluka, Buchs, Switzerland.

30 In a second preferred embodiment of the invention, the cytotoxic portion is a dimeric compound with ribonucleolytic activity, such as a ribozyme,

but preferably ribonuclease (RNase). The enzymes of the RNase family are able to degrade single-stranded RNA molecules to smaller polynucleotides and are directly cytotoxic when intracellular. Bovine seminal RNase (BSRNase) has activities in addition to its RNA-degrading activity, namely anti-tumour (Vescia *et al* (1980) *Cancer Res.* **40**, 3740-3744; Vescia & Tramontano (1981) *Mol. Cell. Biochem.* **36**, 125-128); immunosuppressive (Tamburrini *et al* (1990) *Eur. J. Biochem.* **190**, 145-148; activation by interferon- $\gamma$  (Schein *et al* (1990) *Nucl. Acids Res.* **18**, 1057) and anti-spermatogenic (Doital & Matonsek (1973) *J. Reprod. Fertil.* **33**, 263-274). BSRNase is a dimer and forms two unique disulphide bridges across the subunit interface (Piccoli *et al* (1988) *Biochem J.* **253**, 329-336). The cDNA encoding the precursor to BSRNase can be prepared using the methods disclosed by Preub *et al* (1990) *FEBS Lett.* **270**, 229-232.

Of course, for the invention to work the cytotoxic portion may comprise intact BSRNase, or it may comprise a fragment or fragments of BSRNase retaining at least the active site and subunit-binding sites.

It is further preferred if the fusion with the RNase comprises the sequence KDEL (SEQ ID No 29) at, or near to, the C-terminus of the protein.

It is still further preferred if a linker sequence is present at the N-terminus of the RNase to allow the N-terminus to be more flexible and increase the likelihood of dimer formation.

Preferably, a disulphide-loop-containing sequence which allows an RNase to be linked to a ScFv via a disulphide bond is present in a fusion protein.

In one embodiment the invention, the cytotoxic portion is a compound

with DNA endonucleolytic activity such as copper-phenanthroline adducts but preferably is a DNA endonuclease, for example deoxyribonuclease-I (DNase-I), which is an endonuclease which cleaves double-stranded DNA to yield 5' phosphorylated polynucleotides. It does not cut all DNA sites with the same frequency as it is affected by the local structure of the DNA (specifically, the size of the minor groove).

Alternatively, the DNA endonuclease could be a type II restriction endonuclease. Type II restriction endonucleases are enzymes isolated from microorganisms, usually bacteria, which cleave double-stranded DNA at specific sequences. Typically, the type II restriction endonucleases recognise palindromic sequences in DNA and cleave both strands of the DNA within or adjacent the recognition site. Type II restriction enzymes are dimers of identical subunits, and, for example, *EcoRI* is a homodimer of 31 kDa subunits which recognises the sequence 5'-GAATTC-3'.

Other type II restriction enzymes recognise different hexonucleotide sequences, for example *BamHI* recognises 5'-GGATCC-3', *HindIII* recognises 5'-AAGCTT-3'. In addition, type II restriction enzymes which recognise different numbers of bases are known, for example, *MspI* recognises 5'-CCGG-3', *Sau3AI* recognises 5'-GATC-3', *Hinfi* recognises 5'-GANTC-3' and *NorI* recognises 5'-GCGGCCGC-3'. Of course, the fewer specific bases in the recognition sequence, the more likely that any DNA molecule will be cleaved by the cognate type II endonuclease.

The gene for the bovine DNase I has been chemically synthesized and expressed in *E. coli* (Worrall & Connolly (1990) *J. Biol. Chem.* **265**, 21889-21895. The gene for the human enzyme has been cloned, from a human pancreatic cDNA library constructed in  $\lambda$ gt10 and the enzyme has

been expressed in human cell culture and used in the relief of cystic fibrosis symptoms, by reducing the viscosity of sputum, by degrading the viscous DNA (Shak *et al* (1990) *Proc. Natl. Acad. Sci.* **87**, 9188-9192; Hubbard *et al* (1992) *N. Engl. J. Med.* **326**, 812-815). All the enzymes  
5 are compact, monomeric proteins of about 29 kDa (260 amino acids); when glycosylated the human enzyme is about 35 kDa. It is dependent on divalent cations for activity ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ). The human enzyme is about 75% identical to the bovine enzyme, at the amino acid sequence level. The synthetic gene encoding the bovine DNase-I can be prepared using the  
10 methods disclosed by Worrall & Connolly (1990) *loc. cit.*

The enzyme from bovine pancreas has been purified and crystallized, and a high resolution structure determined at 2Å (Suck & Oefner (1986) *J. Mol. Biol.* **192**, 605-632).

15 One aspect of the invention is the introduction into the targeted cell of the DNase I enzyme. During stages of mitosis, when the nuclear membrane is dissolved, the chromosomal DNA of the said targeted cell will be susceptible to nuclease attack. In this embodiment of the invention  
20 DNase I will be particularly cytotoxic to rapidly dividing cells, such as tumour cells.

A further aspect of the invention is the incorporation into the compound of the invention a nuclear localisation sequence from the SV40 large T  
25 antigen (Kalderon *et al* (1984) *Cell* **39**, 499-509). The said nuclear localisation sequence is PKKKRKV (SEQ ID No.1), or analogues thereof, and a DNA fragment encoding the said sequence, or analogues thereof, may or may not be incorporated into the gene expressing the compound of the invention containing DNase I as the second portion.

Inclusion of the said nuclear localisation sequence will allow the compound of the invention to gain access to the chromosomal DNA during the periods of the cell cycle when the nuclear membrane is intact, as the nuclear pores are permeable to large macromolecules incorporating the  
5 said nuclear localisation sequence, or analogues thereof.

For the invention to work, of course, the cytotoxic portion may comprise a fragment of RNase or of DNA endonuclease which retain their enzymatic activity, such as the active site, and in the case of the dimeric  
10 RNase, and restriction endonuclease, their subunit binding site.

A further aspect of the invention is that the RNase and the DNase are of mammalian, preferably human, origin. The use of the said mammalian proteins as the second, functional portion of the compound of the  
15 invention is advantageous as such compounds are less likely to give rise to undesirable immune reactions.

Many target cell-specific molecules are known, such as those disclosed in Table 1, which are not joined to a further directly or indirectly cytotoxic  
20 portion, but may nevertheless be useful in directing cytotoxic agents to a target cell.

Thus in a further aspect of the invention a compound comprises a mediator portion and a directly or indirectly cytotoxic portion. The  
25 mediator may recognise the native target cell-specific molecule, but it is preferable for the mediator to recognise a derivative of the said molecule.

In the case of antibodies, the native target cell-specific molecule may be recognised by the mediator via its Fc portion.

The said derivative may be made by joining a moiety, such as a small molecule, for example a hapten, to the said molecule, and may be recognised, if the mediator is, for example, an antibody or fragment thereof.

5

The advantage in using this method is that the same moiety may be joined to all types of target cell-specific molecules, and then only one compound, comprising a mediator which recognises the said moiety and a directly or indirectly cytotoxic portion, may be used to deliver the cytotoxic agent to the target cell.

10

In one embodiment of the invention the mediator is ScFV<sub>NP</sub>, and the moiety recognised by the said ScFV<sub>NP</sub> is the hapten 4-hydroxy-3-nitrophenylacetic acid (NP) or 4-hydroxy-3-iodophenylacetic acid, and the target cell-specific molecule is an antibody.

15

Other haptens are suitable as are other molecules, such as peptides, that can be recognised by the mediator. Conveniently the peptide is the core mucin peptide.

20

Before such molecules can be regarded as suitable candidates, there is a requirement that cell specificity be demonstrated and a further requirement that this specificity be shown to be conferred only by the combination of the interaction of the primary targeting antibody with target, and the interaction of the second step reagent (in this case the ScFv) with the primary antibody. To this end, the primary antibody needs to be recognised specifically by the mediator, and therefore requires stable modifications that will distinguish it from native antibodies. Multiple derivatisation of the primary antibody with a hapten fulfils this demand, and has the further advantage of amplification, providing an array of

25

30

secondary targets for the mediator.

Of course, other mediators such as Fab, F(ab')<sub>2</sub>, dAbs or other antibody fragments may be used. The mediator may also recognise the moiety in  
5 a non-immune sense, such as in biotin-streptavidin recognition. It is preferred if the moiety recognised is a small molecule, but the moiety may also be a polypeptide, peptide, oligosaccharide or the like.

The murine immune response to the haptens 4-hydroxy-3-nitrophenylacetic  
10 acid (NP) and 4-hydroxy-3-iodo-5-nitrophenylacetic acid (NIP) is dominated by well characterised V<sub>H</sub> domains and a  $\lambda_1$  light chain (Kabat *et al* (1987) *Sequences of proteins of immunological interest*, US Department of Health and Human Services, Public Health Services, National Institutes of Health). NP-specific V<sub>H</sub> domains have been used in  
15 the construction of recombinant antibodies (Neuberger *et al* (1984) *Nature* 312, 604-608, Casadei *et al* (1990) *Proc. Natl. Acad. Sci. USA* 87, 2047-2051). The hapten itself is well studied and of some immunological interest (Brownstone *et al* (1966) *loc. cit.*) and is also available commercially in a variety of chemical forms. It is relatively simple to  
20 conjugate NP or NIP to other proteins including antibodies.

We describe in the Examples the construction and characterisation of a ScFv with an affinity in the range of  $1-3 \times 10^8 \text{ M}^{-1}$  at pH 7.4 for NIP conjugated to BSA, sufficiently high that the molecule is suitable as a  
25 second step targeting reagent. Derivatisation with hapten resulted in reduction in immunoreactivity of the primary antibody, but even under these adverse circumstances the hapten-conjugated antibody was still capable of delivering ScFv<sub>NP</sub> specifically to cells. Since about forty hapten molecules were conjugated, on average, to each mAb molecule, there is  
30 still a potential 40-fold amplification provided. The specificity of targeting

is governed by the interactions of primary antibody with target, and the ScFv<sub>NP</sub> with derivatised primary antibody, since the ScFv does not bind cells and non-derivatised antibodies bound at cells cannot capture the ScFv. The ScFv described here can therefore be considered as a universal  
5 agent for delivery of drugs or radionuclides or other cytotoxic agents to any cell type for which a previously characterised antibody exists.

In this aspect of the invention, the cytotoxic portion joined to the mediator portion may be a drug, pro-drug, radionuclide, protein including an  
10 enzyme, antibody or any other therapeutically useful reagent.

Thus, the drug may be a cytotoxic chemical compound such as methotrexate, adriamycin, vinca alkaloids (vincristine, vinblastine, etoposide), daunorubicin or other intercalating agents. The enzyme, or  
15 enzymatic portion thereof, may be directly cytotoxic, such as DNaseI or RNase, or indirectly cytotoxic such as an enzyme which converts a substantially non-toxic pro-drug into a toxic form. The protein may be ricin. The cytotoxic portion may comprise a highly radioactive atom, such as iodine-131, rhenium-186, rhenium-188 or yttrium-90, which emits  
20 enough energy to destroy neighbouring cells.

An indirectly cytotoxic portion may be a small-molecule binding site wherein the said small-molecule is capable of being conjugated to any from the following cytotoxic compounds: radioactive compound; drug;  
25 pro-drug; radionuclide; protein including enzyme; antibody; or toxin:

We hereby disclose the principle that ScFvs are suitable for indirect targeting. Moderating the degree of derivatisation of the primary antibody will reduce the loss of immunoreactivity of the primary antibody whilst  
30 still maintaining an array of secondary targets for the hapten-specific



ScFv.

In a further embodiment, the cytotoxic portion of the compound comprises at least the biotin-binding portion of streptavidin as disclosed in Example

5 4.

The compounds of the invention are administered in any suitable way, usually parenterally, for example intravenously, intraperitoneally or, preferably (for bladder cancer), intravesically (ie into the bladder), in  
10 standard sterile, non-pyrogenic formulations of diluents and carriers, for example isotonic saline (when administered intravenously).

A further aspect of the invention provides a method of delivery of the compound of the invention which contains a binding site for a small  
15 molecule, and the administration of the said small molecule conjugated with any from the following: drug, pro-drug, radionuclide, enzyme, antibody or any other therapeutically useful reagent, to give the "small molecule conjugate".

20 Once the compound has bound to the target cells and been cleared from the bloodstream (if necessary), which typically takes a day or so, the small molecule conjugate is administered, usually as a single infused dose. If needed, because the compound of the invention may be immunogenic, cyclosporin or some other immunosuppressant can be administered to  
25 provide a longer period for treatment but usually this will not be necessary.

The timing between administrations of the compound and the small molecule conjugate may be optimised in a non-inventive way since target  
30 cell/normal tissue ratios of conjugate (at least following intravenous

delivery) are highest after about 4-6 days, whereas at this time the absolute amount of antibody bound to the tumour, in terms of percent of injected dose per gram, is lower than at earlier times. Therefore, the optimum interval between administration of the conjugate and the small molecule conjugate will be a compromise between peak target concentration of enzyme and the best distribution ratio between target and normal tissues.

The dosage of the small molecule conjugate will be chosen by the physician according to the usual criteria. The dosage of the compound of the invention will similarly be chosen according to normal criteria, and, in the case of tumour treatment, particularly with reference to the type, stage and location of the tumour and the weight of the patient. The duration of treatment will depend in part upon the rapidity and extent of any immune reaction to the antibody or cytotoxic component of the compound.

A further aspect of the invention provides a method of delivery of the target cell-specific molecule and a compound of the invention which contains a mediator portion. Once the target cell-specific molecule has bound to the target cells and been cleared from the bloodstream (if necessary), which typically takes a day or so, the compound comprising a mediator portion is administered in any suitable way.

If the cytotoxic portion, joined to the mediator portion, contains a binding site for a small molecule, then, once the mediator-containing compound has bound to the target cell-specific molecule at the site of the target cell, and has been cleared from the bloodstream (if necessary), the said small molecule conjugate is administered as described *supra*.

The compounds of the invention either in themselves, or together with a target cell-specific molecule or additionally together with an appropriate toxic agent, capable of binding to the small molecule-binding site of the compound, are in principle suitable for the destruction of cells in any tumour or other defined class of cells selectively exhibiting a recognisable (surface) entity. The compounds are principally intended for human use but could be used for treating other mammals including dogs, cats, cattle, horses, pigs and sheep.

- 10 The small molecule conjugate, when used in combination with a compound for diagnosis, usually comprises a radioactive atom for scintigraphic studies, for example technetium 99m ( $^{99m}\text{Tc}$ ) or iodine-123 ( $^{123}\text{I}$ ), or a spin label for nuclear magnetic resonance (nmr) imaging (also known as magnetic resonance imaging, mri), such as iodine-123 again, iodine-131, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese or iron.

- When used in combination with a compound for selective destruction of the tumour, the small molecule conjugate may comprise a highly radioactive atom, such as iodine-131, rhenium-186, rhenium-188 or yttrium-90, which emits enough energy to destroy neighbouring cells, or a cytotoxic chemical compound such as methotrexate, adriamycin, vinca alkaloids (vincristine, vinblastine, etoposide), daunorubicin and other intercalating agents or (preferably) an enzyme or enzymatic portion thereof which converts a non-toxic pro-drug into a toxic form. In the latter case, the compound of the invention is administered and, once there is an optimum balance between (i) the tumour to normal cell ratio of compound and (ii) the absolute level of compound associated with the tumour, the pro-drug is administered either systemically (eg intravenously) or intravesically, into the bladder. The enzyme/pro-drug systems of

Bagshawe and his co-workers may be used (*loc. cit.*) or the antibody-alkaline phosphatase conjugates, followed by etoposite phosphate (*loc. cit.*) or, more preferably, the cyanide-liberating systems described by Epenetos (*loc. cit.*).

5

The compounds of the invention, together with an appropriate small molecule conjugated to a readily-detectable reagent such as a radionuclide; fluorescent molecule; or enzyme are in principle suited for the recognition of antigens in other situations. These include immunoblotting procedures, such as the well-known Western blot (Towbin *et al* (1979) *Proc. Natl. Acad. Sci. USA* 76, 4350-4354); assays such as the enzyme-linked immunosorbent assay (ELISA); and *in situ* hybridisation experiments in which the presence of antigens within fixed cells is detected.

10

15 In a further embodiment of the invention, a compound comprising an oligomeric complex of at least two molecules each comprising a target cell-specific portion and a further portion wherein the molecules are complexed to one another via their further portions is useful in agglutinating cells. In a preferred embodiment the target cell-specific  
20 portion of the compound of the invention recognises particular blood group antigens displayed on the surface of the erythrocyte, and because of the multivalent binding nature of the compound, the addition of the compound to blood may lead to haemagglutination. Thus, in this embodiment the compounds may be specific to particular antigens within  
25 the ABO, Rhesus, Kell, or any other blood group systems, and the compound of the invention may find uses in blood group typing or other areas of tissue typing.

Antibodies, including monoclonal antibodies, are known which react with  
30 most of the aforementioned blood group antigens, and it is well within the

scope of a person skilled in the art to derive, for example, ScFvs from such antibodies for use in the invention.

5 The invention will now be described in detail with reference to the following figures and examples wherein:

Figure 1 shows a diagrammatic representation of compounds of the invention.

10 Figure 2 shows the construction of plasmids expressing ScFv<sub>NP</sub>.

Figure 3 shows oligonucleotide primers used in the polymerase chain reaction to amplify various fragments of the ScFv coding region.

15 Figure 4 shows the nucleotide sequence (SEQ ID No 2) (and encoded protein sequence (SEQ ID No 3)) between the *Hind*III and *Eco*RI sites of pRAS107 and pRAS111.

20 Figure 5 shows the binding of a soluble protein expressed from pRAS111 to NIP<sub>15</sub>-BSA.

Figure 6 shows that a soluble protein expressed from pRAS111 and which binds NIP<sub>15</sub>-BSA can be competed by NIP<sub>15</sub>-BSA.

25 Figure 7 shows the construction of plasmids expressing ScFv-streptavidin fusions *in vitro*.

Figure 8 shows the construction of plasmids for the expression of ScFv-streptavidin fusions in *E. coli*.

Figure 9 shows the nucleotide sequence (SEQ ID No 4) (and deduced amino acid sequence (SEQ ID No 5)) between the *HindIII* and *EcoRI* sites of pRAS108 and pRAS112.

- 5 Figure 10 shows the nucleotide sequence (SEQ ID No 6) (and deduced amino acid sequence (SEQ ID No 7)) between the *HindIII* and *EcoRI* sites of pRAS109 and pRAS113.

- 10 Figure 11 shows the nucleotide sequence (SEQ ID No 8) (and deduced amino acid sequence (SEQ ID No 9)) between the *HindIII* and *EcoRI* sites of pRAS110 and pRAS114.

Figure 12 shows the detection of soluble pRAS112-encoded protein (full length ScFv<sub>NP</sub>-streptavidin monomer) in bacterial supernatants.

15

Figure 13 shows that pRAS112-encoded protein binds to NIP<sub>15</sub>-BSA, but not to lysozyme.

20

Figure 14 shows that concentrated pRAS112-encoded protein binds iminobiotin-Sepharose at pH 11 in contrast to parental ScFv<sub>NP</sub> protein that does not.

25

Figure 15 shows a diagrammatic representation of pRAS112-encoded protein.

Figure 16 shows the construction of plasmids expressing ScFv-BSRNase fusion molecules.

30

Figure 17 shows a diagrammatic representation of a ScFv-BSRNase heterodimer.

Figure 18 shows the construction of plasmids expressing ScFv-DNaseI fusion molecules.

Figure 19 shows the purification of pRAS111 ScFv<sub>NP</sub> protein.

5

Figure 20 shows indirect targeting of pRAS111 ScFv<sub>NP</sub>.

Figure 21 shows the nucleotide sequence (SEQ ID No 10) of the ScFv-BSRNase fusion (anti-4-OH-nitrophenacetyl antibody) that has been  
10 inserted between the *Hind*III and *Eco*RI sites of plasmid pSP71.

Figure 22 shows the nucleotide sequence (SEQ ID No 11) of the ScFv-BSRNase fusion (H17-BSRNase; anti-human placental alkaline  
phosphatase antibody; H17E2) that has been inserted between the *Hind*III  
15 and *Eco*RI sites of plasmid pSP71.

Figure 23 shows the nucleotide sequence (SEQ ID No 12) of the ScFv-BSRNase fusion (anti-lysozyme antibody) that has been inserted between  
the *Hind*III and *Eco*RI sites of a plasmid pUC18.

20

Figure 24 shows the nucleotide sequence (SEQ ID No 13) of the ScFv-DNaseI fusion (anti-4-OH nitrophenacetyl antibody) that has been inserted  
between the *Hind*III and *Bgl*II sites of plasmid pSP71.

25 Figure 25 shows the nucleotide sequence (SEQ ID No 14) of the ScFv-DNaseI fusion (anti-human placental alkaline phosphatase antibody;  
H17E2) that has been inserted between the *Hind*III and *Bgl*II sites of  
plasmid pSP71.

30 Figure 26 shows the nucleotide sequence (SEQ ID No 15) of the ScFv-

DNaseI fusion (anti-lysozyme antibody) that has been inserted between the *HindIII* and *BglII* sites of plasmid pUC18.

Figure 27 shows the results of cell-killing experiments using HEp2 cells and the fusion protein H17-DT-BSR, H17-DT-BSR/KDEL and H17-DT-BSR/KDELINK.

Figure 28 is a schematic diagram of the H17E2 scFv-seminal RNase fusion proteins. The plasmid which express them are named in parentheses.

Figure 29 shows the nucleotide sequence (SEQ ID No 24) encoding the H17E2 scFv-diphtheria toxin disulphide loop-BSRNase (H17-Dip. Tox.-BSRNase).

Figure 30 shows the nucleotide sequence (SEQ ID No 25) encoding the H17E2 scFv-diphtheria toxin disulphide loop-BSRNase-KDEL (H17-Dip. Tox.-BSRNase KDEL).

Figure 31 shows the nucleotide sequence (SEQ ID No 26) encoding the H17E2 ScFv diphtheria toxin disulphide loop-Linker-BSRNase-KDEL (H17-Dip. Tox.-link-BSRNase KDEL).

Figure 32 shows the nucleotide sequence (SEQ ID No 27) encoding the H17E2 ScFv-Linker-BSRNase-KDEL (H17-LBSRNase-KDEL).

Figure 33 shows the nucleotide sequence (SEQ ID No 28) encoding the H17E2 ScFv-BSRNase KDEL).

Figure 34 shows the elution of pRAS111 and pRAS112 proteins from NP-



sepharose with 50 mM glycine HCl, pH2.2.

**Example 1: Construction of a single-chain Fv (ScFv) reactive against the hapten NP (4-OH nitrophenacetyl)**

5

**Plasmid constructions**

Plasmids are shown in Figure 2. Filled circles represent promoters:  $P_{lac}$ , *lac* promoter of pUC plasmids;  $P_{SP6}$ , SP6 promoter;  $P_{T7}$ , T7 promoter.  
 10 Open boxes represent fused gene portions: *pelB*, the signal sequence derived from the pectate lyase B gene of *Erwinia caratovora*;  $(G_4S)_3$ , flexible oligopeptide linker comprising three tandem repeats of N-GlyGlyGlyGlySer-C (SEQ ID No 16); *myc*, a small immunogenic tag derived from *c-myc*. Restriction enzyme sites: B, *Bam*HI; Bs, *Bst*EII; b, *Bgl*II; C, *Cla*I; E, *Eco*RI; H, *Hind*III; K, *Kpn*I; P, *Pst*I; Sp, *Sph*I; Ss, *Sst*I; X, *Xho*I.

Plasmid pSWsFvD1.3myc (McCafferty *et al* (1990) *Nature* 348, 552-554) encodes a single-chain Fv reactive against hen egg lysozyme, and which  
 20 comprises  $VH_{D1.3}$  and  $V_{K_{D1.3}}$  domains linked by a flexible oligopeptide,  $(G_4S)_3$ , under the transcriptional control of the *lac* promoter of *E. coli*. The region encoding  $V_{K_{D1.3}}$  was replaced by one encoding  $V_{\lambda}$  in the following manner. The segment encoding  $VH_{D1.3}(G_4S)_3$  was subjected to polymerase chain reaction (PCR) mediated amplification using  
 25 oligonucleotide primers VHBACK2 (SEQ ID No 17) and BAMLINKERFOR (SEQ ID No 18) (Figure 3). Primer BAMLINKERFOR directs the incorporation of a *Bam*HI site that also encodes the two carboxy-terminal amino acids of the flexible oligopeptide linking the two V domains.

A  $V\lambda$  gene segment was amplified from chromosomal DNA of plasmacytoma J558L using primer pair BAMV $\lambda$ BACK (SEQ ID No 19) and ECOV $\lambda$ FOR (SEQ ID No 20). The former directs the incorporation of a *Bam*HI site at the 5' end of the gene; the latter two stop codons and  
5 *Xho*I and *Eco*RI sites at the 3' end of the gene.

The two amplified products were used to replace the *Pst*I-*Eco*RI fragment of plasmid pRAS103 to generate plasmid pRAS106 which encodes a ScFv protein comprising  $VH_{D1.3}(G_4S)_3V\lambda_{J558L}$  under the transcriptional control  
10 of the SP6 promoter.

The *Pst*I-*Bst*EII fragment of pRAS106 was replaced with a *Pst*I-*Bst*EII fragment encoding  $VH_{NP}$  amplified from plasmid pRAS49 (Spooner and Lord (1991) *loc. cit.*) using primers VHBAC3 (SEQ ID No 21) and  
15 VH1FOR-2 (SEQ ID No 22) to generate plasmid pRAS107. This bears a  $VH_{NP}(G_4S)_3V\lambda_{J558L}$  ScFv under the transcriptional control of the SP6 promoter, and is intended purely for expression in *in vitro* systems.

Plasmid pRAS111 bears the ScFv of pRAS107, but under T7 promoter  
20 control, and is suitable for expression in both *in vitro* systems and bacterial systems.

The nucleotide sequence (and deduced amino-acid sequence) between the *Hind*III and *Eco*RI sites of plasmids pRAS107 and pRAS111 are given in  
25 Figure 4.

Table 3: Plasmids used

Plasmid	Relevant characteristics	Source or reference
pSWsFvD1.3myc	Anti-lysozyme ScFv, $VH_{D1.3}(G_4S)_3Vk_{D1.3}$	McCafferty <i>et al</i> (1990) <i>loc. cit.</i>
pRAS103	Anti-lysozyme ScFv-ricin A chain fusion, <i>lac</i> promoter	Spooner <i>et al</i> (1992) pp 7-15 in <i>Monoclonal Antibodies 2; Applications in Clinical Oncology</i> (Epenetos, A.A., Ed), Chapman & Hall
pRAS106	$VH_{D1.3}(G_4S)_3V\lambda_{J558L}$ , SP6 promoter	This application
pRAS49	Anti-NP antibody H chain-ricin A chain fusion, IgH promoter	Spooner and Lord (1991) <i>loc. cit.</i>
pRAS107	$VH_{NP}(G_4S)_3V\lambda_{J558L}$ , SP6 promoter	This application
pRAS111	$VH_{NP}(G_4S)_3V\lambda_{J558L}$ , T7 promoter	This application

#### 10 Growth of plasmacytoma J558L and DNA preparation

Mouse plasmacytoma J558L cells were grown in Dulbecco's modified Eagle's medium supplemented with 10% foetal calf serum. Cells were washed twice in standard phosphate-buffered saline pH 7.4 (PBS) and high molecular weight DNA was prepared by addition, with gentle vortexing, of  $10^6$  cells suspended in 100  $\mu$ l PBS to 2.5 ml 10 mM Tris-HCl 1 mM EDTA pH 8.0 containing 0.02% (w/v) SDS. After adding Proteinase K to 1 mg.ml<sup>-1</sup>, incubation (3h, 50°C) and two phenol/chloroform extractions, DNA was precipitated with ethanol, and dissolved overnight at 4°C in 1 ml 10 mM Tris-HCl 1 mM EDTA pH 8.

### Polymerase chain reaction

Plasmid or chromosomal DNA (100 ng) was subjected to 24 rounds of PCR-mediated amplification (94°C, 1 min; 65°C, 1.5 min; and 72°C, 2 min) in 50  $\mu$ l reaction volumes containing 25 pmol of each appropriate oligonucleotide primer, 250  $\mu$ M of each dNTP, 67 mM Tris-HCl (pH 8.8), 17 mM  $(\text{NH}_4)_2\text{SO}_4$ , 1.5-6 mM  $\text{MgCl}_2$ , 200  $\text{mg}\cdot\text{ml}^{-1}$  gelatin and 5 units of Taq polymerase (Cetus) overlaid with 25  $\mu$ l paraffin oil. Amplified DNA was extracted once with phenol/chloroform and precipitated with ethanol before use.

### Bacterial expression of pRAS111 protein

*E. coli* K12 JM109(DE3), a JM109 derivative with a chromosomal insertion of T7 polymerase under *lac* transcriptional control, was transformed with plasmid pRAS111. Cells were grown to a density of  $10^7$   $\text{ml}^{-1}$  and expression of pRAS111 protein was induced by induction of T7 polymerase with 100 nM IPTG. A 31 kDa protein accumulates in the cells in sufficient quantity for provisional identification by Coomassie staining of cell extracts. The identity is confirmed by Western Blotting, probing with biotinylated goat anti-mouse lambda ( $\text{G}\alpha\text{m}\lambda$ ) antiserum.

In addition, *E. coli* K12 BL21 (DE3), a derivative of BL21 with a single chromosomal copy of T7 RNA polymerase under *lacUV5* promoter control (Studier and Moffatt (1986) *J. Mol. Biol.* 189, 113-130) was transformed with plasmid pRAS111. Cultures (400 ml) were grown at 37°C or at room temperature in minimal salts medium supplemented with 100  $\mu\text{g}\cdot\text{ml}^{-1}$  ampicillin and 1% glucose or in L-broth supplemented with 100  $\mu\text{g}\cdot\text{ml}^{-1}$  ampicillin, to a density of  $10^7$   $\text{cells}\cdot\text{ml}^{-1}$ . Expression of pRAS111 ScFv protein was achieved by induction of T7 polymerase with

100 nM IPTG. After induction, cells were grown for 24 h to permit accumulation of pRAS111 ScFv protein in the growth medium.

### Biological activity and affinity purification of pRAS111 protein

5

Filtered bacterial supernatants were applied to wells of a 96-well plate previously coated with 10 mg.ml<sup>-1</sup>NIP<sub>15</sub>-BSA or 300 mg.ml<sup>-1</sup> hen egg lysozyme, and bound protein was detected by serial incubation with biotinylated G $\alpha$ m $\lambda$  antiserum and HRPO-streptavidin conjugate. Colour changes were generated by incubation with ABTS and were monitored at 405 nm.

10

A soluble protein present in the growth medium of JM109 (DE3)/pRAS111 cultures, but not in cultures of JM109 (DE3), binds NIP<sub>15</sub>-BSA, but not lysozyme (Figure 5). Filtered bacterial growth medium recovered after induction of pRAS111 protein was applied to wells of an ELISA plate coated with 10  $\mu$ g.ml<sup>-1</sup> NIP<sub>15</sub>-BSA (⊙) or 300  $\mu$ g.ml<sup>-1</sup> hen egg lysozyme (♦). Bound protein was detected by serial incubation with biotinylated G $\alpha$ m $\lambda$  (Goat anti-mouse lambda light chain) antisera and horseradish peroxidase conjugated streptavidin diluted in blocking buffer, and colour changes generated by addition of ABTS were monitored at 405 nm. A soluble protein present in the growth medium of JM109(DE3)/pRAS111 cultures, but not in cultures of JM109(DE3), binds NIP<sub>15</sub>-BSA, can be competed with NIP<sub>15</sub>-BSA (Figure 6). ScFv protein was allowed to bind ELISA wells coated with 10  $\mu$ g.ml<sup>-1</sup> NIP<sub>15</sub>-BSA in the absence of competing hapten, or in the presence of 0.010  $\mu$ g.ml<sup>-1</sup> (◇), 0.019  $\mu$ g.ml<sup>-1</sup> (X), 0.039  $\mu$ g.ml<sup>-1</sup> (-), 0.078  $\mu$ g.ml<sup>-1</sup> (+), 0.156  $\mu$ g.ml<sup>-1</sup> (♦), 0.313  $\mu$ g.ml<sup>-1</sup> (✦), 0.625  $\mu$ g.ml<sup>-1</sup> (Δ), 1.25  $\mu$ g.ml<sup>-1</sup> (■), 2.5  $\mu$ g.ml<sup>-1</sup> (□), 5  $\mu$ g.ml<sup>-1</sup> ( ) or 10  $\mu$ g.ml<sup>-1</sup> (○) competing hapten. Bound protein

20

25

was detected by serial incubation with biotinylated G $\alpha$ m $\lambda$  antisera and horseradish peroxidase conjugated streptavidin, and colour changes generated by addition of ABTS were monitored at 405 nm. The ScFv encoded by pRAS111 was found to have a binding affinity for NP of  $K_d$   
5 =  $4 \times 10^{-9}$  M. Since bivalency of an antibody commonly provides an extra three orders of magnitude of binding ability, then an avidity of at least  $10^{-12}$  M would be predicted for bivalent molecules derived from ScFvNP.

As an alternative, growth medium, filtered through 0.2  $\mu$ m nitrocellulose  
10 filters to remove cells and particulates, was adjusted to 80% saturation with solid ammonium sulphate at 4°C. After incubation (4°C, 1 h) treated medium was centrifuged (10,000 x g, 30 min) to pellet insoluble proteins. Pellets were taken up in 20 ml PBS and were dialysed exhaustively against PBS at 4°C. Insoluble material after dialysis was  
15 removed by brief centrifugation and the remainder was adjusted to 40 ml final volume with PBS, to 0.02% with sodium azide and was applied slowly (2 ml h<sup>-1</sup>) to a 2 ml NP-Sepharose column at room temperature. After washing with 50 column volumes of PBS containing 0.02% sodium azide (PBS/azide), bound proteins were eluted with 50 mM glycine-HCl  
20 pH 2.2 and fractions (2 ml) were immediately adjusted by addition of 200  $\mu$ l 2M unbuffered Tris base. Fractions containing ScFv protein were pooled, dialysed against PBS and concentrated using Macrosep (Amicon) concentrators with a 10 kDa cut-off. Yields were estimated by Bio-Rad protein assay, using rabbit IgG as a reference, and by absorbance at 280  
25 nm assuming  $A_{280}=1$  for 1.4 mg.ml<sup>-1</sup> solution.

Soluble NIP-binding activity was detected by ELISA analysis of bacterial growth medium after induction, and could be concentrated by ammonium sulphate precipitation and purified by affinity chromatography on NP-  
30 Sepharose (Fig 19) so no attempt was made to recover pRAS111 ScFv

protein present in cell pellets. Yields of pRAS111 ScFv from growth medium were not greatly different when induced at room temperature or 37°C. Induction of expression was efficient in minimal salts medium and not discernible in rich broth; however, little difference was noted in the final yields. The most important factor found here was the bacterial strain, with yields of  $\sim 1.3 \text{ mg l}^{-1}$  pRAS111 ScFv protein recovered from cultures of BL21(DE3)/pRAS111, approximately ten-fold greater than those obtained from cultures of JM109(DE3)/pRAS111.

#### 10 Specificity of pRAS111 protein

The screening agent used here, biotinylated G $\alpha$ m $\lambda$  antiserum, also detects the  $\lambda$ 1 light chain of anti-NP/NIP antibodies. It is therefore not possible to demonstrate specificity of pRAS111 ScFv protein for NP or NIP by competition with anti-NP/NIP antibodies, but only by its ability to recognise NP/NIP. A soluble protein present in growth medium of JM109 (DE3)/pRAS111 cultures, but not in untransformed cultures of JM109 (DE3), binds NIP<sub>16</sub>-BSA, but not lysozyme, and can be competed with NIP<sub>16</sub>-BSA (Fig 5). This activity can be retained on NP-Sepharose columns, from where it can be eluted. In addition, targeting studies demonstrate no cross-reaction with BSA, PEPY2-BSA, antibody or mammalian cells.

#### Affinity determinations

Results of affinity determinations using ELISA-based techniques are given in Table 4. Affinity of pRAS111 ScFv for NIP was estimated firstly by adapting the method of Mariani *et al* (1987) *Molec. Immunol.* **24**, 297-303), determining the concentration of total added antibody giving half-maximal binding ( $C_{50}$ ) assuming  $C_{50} = 1/K_{app}$ , where  $K_{app}$  is the apparent

affinity constant. This approximation only holds true if the number of available binding sites per well is sufficiently low that their contribution is insignificant. Determinations of  $K_{app}$  should approach  $K_{actual}$  as the amount of antigen per well is reduced. Table 4 shows that a point is  
 5 reached where similar values of  $K_{app}$  are derived ( $K = 2-3 \times 10^8 \text{ M}^{-1}$ ), representing the closest approximation that can be made using this method.

To confirm the accuracy of this approach, similar estimations of  $K$  were made using the method of Hogg *et al* (1987) *Molec. Immunol.* 24, 797-  
 10 801) in the absence of competing antigen, by calculating the slope of the linear portion of a plot of  $A_{450}/[\text{ScFv}_{NP}] \text{ v } A_{450}$ , where  $A_{450}/[\text{ScFv}_{NP}] = fKn - fK(A_{450})$ ,  $A_{450}$  is the absorbance at 450 nm,  $[\text{ScFv}_{NP}]$  is the concentration of added  $\text{ScFv}_{NP}$ ,  $n$  is the concentration of available binding sites and  $f$  is the valency of the  $\text{ScFv}_{NP}$  for NIP. A value of 1 was  
 15 assigned to  $f$ .

Table 4: Affinity determinations of pRAS111  $\text{ScFv}_{NP}$  protein

Antigen coat	concn (mg ml <sup>-1</sup> ) of coating buffer	K (M <sup>-1</sup> ) Mariani <i>et al</i> (1987), Hogg <i>et al</i> (1987)	
NIP <sub>16</sub> BSA	5	2.5 ( $\pm$ 0.1) $\times 10^9$	1.6 ( $\pm$ 0.1) $\times 10^9$
NIP <sub>16</sub> BSA	1	8.2 ( $\pm$ 1.5) $\times 10^8$	8.1 ( $\pm$ 0.7) $\times 10^8$
NIP <sub>4</sub> BSA	10	2.9 ( $\pm$ 0.3) $\times 10^8$	1.2 ( $\pm$ 0.2) $\times 10^8$
NIP <sub>4</sub> BSA	5	2.5 ( $\pm$ 0.5) $\times 10^8$	1.8 ( $\pm$ 0.1) $\times 10^8$

#### Preparation of NP-Sepharose

Sepharose support (20 ml) with an amine function (Affigel 102, Biorad) was washed and suspended by addition of 20 ml 40 mM triethylamine.



To this was added 430 mg NP-cap-OSu (Cambridge Research Biochemicals) dissolved in 1 ml dimethylformamide (DMF). After mixing by gentle inversion (2h, room temperature) and extensive washing in water and then PBS, NP-Sepharose was equilibrated in PBS/azide and stored in the dark at 4°C.

### Western blots to identify pRAS111 protein

Western blots were performed as previously described (Spooner and Lord (1991) pp 65-77 in *Monoclonal Antibodies; Applications in Clinical Oncology* (Epenetos, A.A., Ed) Chapman and Hall) and pRAS111 ScFv protein was identified by serial incubations in PBS/5% milk powder/0.1% Tween 20 (blocking solution), biotinylated G $\alpha$ m $\lambda$  antisera and streptavidin-HRPO diluted in blocking solution to concentrations recommended by the suppliers. After each incubation, blots were washed 5 times in PBS/0.1% Tween 20. Proteins bound by biotinylated G $\alpha$ m $\lambda$  antisera and streptavidin-HRPO were revealed by incubation with DAB.

### Example 2: Derivatisation of proteins with hapten

20

NIP-cap-OSu (Cambridge Research Biochemicals) was dissolved in dimethylformamide to 20 mg.ml<sup>-1</sup> and added to proteins as below.

NIP-BSA: for low coupling ratio, 80  $\mu$ l 20 mg.ml<sup>-1</sup> NIP-cap-OSu/DMF was added to 1 ml 200 mg.ml<sup>-1</sup> BSA in 10 mM triethylamine. For high coupling ratio, 800  $\mu$ l 20 mg.ml<sup>-1</sup> NIP-cap-OSu/DMF was added to 1 ml 200 mg.ml<sup>-1</sup> BSA in 100 mM triethylamine.

NIP-antibody: 200  $\mu$ l 20 mg.ml<sup>-1</sup> NIP-cap-OSu/DMF was added to 2 ml 2.8 mg.ml<sup>-1</sup> antibody (AUA1 or HMFG1, Unipath) in PBS/40 mM

triethylamine.

After mixing by inversion (2 h, room temperature) and extensive dialysis against PBS, insoluble material was removed by centrifugation. Soluble NIP-BSA was adjusted to 0.02% with sodium azide. Soluble NIP-antibody was sterilised by filtration (0.2  $\mu$ m filter). Haptenated proteins were stored at 4°C in the dark. Protein concentration was estimated by Bio-Rad protein assay.

10 The number of haptens conjugated to each protein molecule was estimated by absorbance at 430 nm according to Brownstone *et al* (1966) *Immunology* 10, 465-479: low coupling ratio NIP-BSA, 3.7 (NIP<sub>4</sub>-BSA); high coupling ratio NIP-BSA, 16.4 (NIP<sub>16</sub>-BSA); NIP-AUA1, 38.3 (NIP<sub>38</sub>-AUA1) and NIP-HMFG1, 35.4 (NIP<sub>35</sub>-HMFG1).

15

### Example 3: Indirect targeting using pRAS111 ScFv protein

The measured affinity of ScFv<sub>NP</sub> or pRAS111 protein is sufficiently high to contemplate cell targeting by a two-step approach. Cells (LOVO and HT29) and peptide (PEPY-BSA) were incubated with AUA1, NIP<sub>38</sub>-AUA1, or NIP<sub>35</sub>-HMFG1, and bound material was detected by incubation with sheep anti mouse antisera (Sh $\alpha$ m) conjugated to HRPO or by serial incubation with biotinylated G $\alpha$ m $\lambda$  and streptavidin-HRPO (Fig 20). LOVO cells, which express AUA1 antigen, can be identified by serial incubation with specific antibody (AUA1) and with Sh $\alpha$ m conjugated to HRPO. Hapten-derivatised NIP<sub>38</sub>-AUA1 displayed a marked reduction in cell-binding ability, with loss of approximately 90% of immunoreactivity. Hapten-conjugated NIP<sub>35</sub>-HMFG1 also bound LOVO cells, reflecting the ability of HMFG1 to bind these cells when presented at high concentration. When pRAS111 ScFv<sub>NP</sub> protein was used as a detection

20

25

30

layer, hapten-derivatised NIP<sub>38</sub>-AUA1 and NIP<sub>35</sub>-HMFG1 were both recognised, but non-hapten-conjugated AUA1 was not. Similar results were obtained with a different cell line, HT29, that also expresses AUA1 antigen.

5

When the specificity of the system was altered completely, a peptide (PEPY2) derived from the protein backbone of polymorphic epithelial mucin identified with NIP<sub>35</sub>-HMFG1 antibody was bound by pRAS111 ScFv protein whilst those incubated with AUA1 and NIP<sub>38</sub>-AUA1 were not.

10

The specificity of pRAS111 ScFv<sub>NP</sub> protein is therefore dependent upon prior targeting with a hapten-derivatised primary targeting vehicle, and the specificity of targeting depends only upon the interaction of primary

15 hapten-conjugated targeting vehicle and the interaction of second step ScFv with the primary targeting vehicle.

20

For ELISAs using fixed mammalian cells, cells were seeded into wells of 96-well microculture plates at  $10^5$  cells.ml<sup>-1</sup> in RPMI supplemented with 10% foetal calf serum and were grown to confluence at 37°C in a 5% CO<sub>2</sub> atmosphere. Cells were washed twice in PBS, were incubated in 0.25% glutaraldehyde in PBS (100 µl per well, room temperature, 15 min) and after a further wash in PBS, were stored at 4°C in PBS/azide.

25

Unbound sites were blocked (30 min, room temperature) using 1% milk powder reconstituted in PBS containing 0.1% Tween 20 (blocking buffer). Antibodies and hapten-conjugated antibodies were applied and were detected by serial incubation with pRAS111 ScFv protein, biotinylated Gαmλ antisera and streptavidin-HRPO or by incubation with horseradish

30 peroxidase conjugated Sheep anti mouse serum, diluted in blocking buffer

to appropriate concentrations. After each incubation, plates were washed 5 times in PBS containing 0.1% Tween 20. Colour changes were generated using ABTS (monitored at 405 nm) or OPD (monitored at 450 nm).

5

The results of indirect targeting of pRAS111 ScFv<sub>NP</sub> are shown in Figure 20.

Binding of AUA1 (open circles), NIP<sub>38</sub>-AUA1 (closed circles) and NIP<sub>35</sub>-HMFG1 (open triangles) to LOVO cells, HT29 cells and to a peptide derived from the mucin backbone conjugated to BSA (PEPY2-BSA). Bound primary antibody was detected using HRPO-conjugated sheep anti-mouse antisera (Sham) or by recognition using pRAS111 ScFv<sub>NP</sub> (ScFv).

#### 15 Example 4: Construction of high avidity ScFv-streptavidin fusion

##### Plasmid constructions

Plasmids for the *in vitro* expression of ScFv-streptavidin fusions are shown in Figure 7. Filled circles represent promoters: P<sub>SP6</sub>, SP6 promoter; P<sub>T7</sub>, T7 promoter. Open boxes represent fused gene portions: *pelB*, the signal sequence derived from the pectate lyase B gene of *Erwinia caratovora*; (G<sub>4</sub>S)<sub>3</sub>, flexible oligopeptide linker comprising three tandem repeats of N-GlyGlyGlyGlySer-C; A-P, a novel flexible oligopeptide linker.

25

Restriction enzyme sites: B, *Bam*HI; Bs, *Bst*EII; b, *Bgl*II; C, *Cla*I; E, *Eco*RI; H, *Hind*III; K, *Kpn*I; P, *Pst*I; Sp, *Sph*I; Ss, *Sst*I; X, *Xho*I; Xb, *Xba*I.

30 Plasmids for the expression of ScFv-streptavidin fusions in *E. coli* are

shown in Figure 8. Filled circles represent promoters:  $P_{SP6}$ , SP6 promoter;  $P_{T7}$ , T7 promoter. Open boxes represent fused gene portions: *pelB*, the signal sequence derived from the pectate lyase B gene of *Erwinia caratovora*;  $(G_4S)_3$ , flexible oligopeptide linker comprising three tandem repeats of N-GlyGlyGlyGlySer-C; A-P, a novel flexible oligopeptide linker. Restriction enzyme sites: B, *Bam*HI; Bs, *Bst*EII; b, *Bgl*II; C, *Cla*I; E, *Eco*RI; H, *Hind*III; K, *Kpn*I; P, *Pst*I; Sp, *Sph*I; Ss, *Sst*I; X, *Xho*I; Xb, *Xba*I.

10. Segments of DNA encoding mature streptavidin monomers or fragments were amplified by PCR and were used to replace the *Xho*I-*Eco*RI fragment of plasmid pRAS107 to generate plasmids pRAS108, pRAS109 and pRAS110, which encode ScFv<sub>NP</sub>-streptavidin fusions under SP6 transcriptional control.

15

Plasmid pRAS108 encodes a ScFv<sub>NP</sub> fused via a novel oligopeptide (APAAAPA (SEQ ID No 23)). Its product is expected to tetramerise via the streptavidin monomer moieties. Mature streptavidin often forms higher order complexes, probably through interaction of the amino-terminal and carboxy-terminal regions which are thought to be flexible extensions. Many commercial preparations lack these, through natural proteolysis, and form stable tetramers. In order to mimic this, two further ScFv<sub>NP</sub>-streptavidin derivatives were made, one borne on plasmid pRAS109 and which lacks the 19 carboxy terminal amino acids of streptavidin, and the other, on plasmid pRAS110, which further lacks the 12 amino-terminal amino acids of streptavidin. Plasmid pRAS110 thus encodes a ScFv<sub>NP</sub> linked to "core" streptavidin monomers, typical of many commercial preparations.

- 30 Plasmids pRAS112, pRAS113 and pRAS114 are derived from plasmids

pRAS108, pRAS109 and pRAS110 respectively, and code for ScFv<sub>NP</sub>-streptavidin fusions under the transcriptional control of the T7 promoter.

The nucleotide sequence (and deduced amino-acid sequence) between the  
5 *Hind*III and *Eco*RI sites of plasmids pRAS108 and pRAS112 are given in  
Figure 9, the sequences of plasmids pRAS109 and pRAS113 in Figure 10  
and those of plasmid pRAS110 and pRAS114 are displayed in Figure 11.

#### Bacterial expression of pRAS112, pRAS113 and pRAS114 proteins

10

In contrast to ScFv<sub>NP</sub>, in the conditions used, proteins encoded by plasmids  
pRAS112, pRAS113 and pRAS114 do not accumulate after induction in  
amounts sufficient for provisional identification by Coomassie staining.  
Western Blotting of cell extracts and culture supernatants, probing with  
15 biotinylated G $\alpha$ m $\lambda$  antiserum and HRPO-streptavidin conjugate or rabbit  
 $\alpha$ -streptavidin (R $\alpha$ S) antiserum and HRPO-donkey  $\alpha$ -rabbit (D $\alpha$ R)  
antiserum allows identification of fusion proteins of expected monomeric  
sizes. Very little ScFv<sub>NP</sub>-core streptavidin accumulates after induction of  
expression of pRAS114 protein.

20

In non-reducing conditions, almost all of the ScFv-streptavidin material  
migrates with sizes corresponding to multimeric forms (at ~90 kDa for  
a dimer and 180 kDa for the tetramer). Note that in the conditions  
employed here, streptavidin itself exists mostly as higher order aggregates.

25

#### Antigen binding

Filtered bacterial supernatants were applied to wells of a 96-well plate  
previously coated with 10  $\mu$ g.ml<sup>-1</sup> NIP<sub>15</sub>-BSA or 300  $\mu$ g.ml<sup>-1</sup> hen egg  
30 lysozyme, and bound protein was detected by serial incubation with

biotinylated  $\text{G}\alpha\text{m}\lambda$  antiserum and HRPO-streptavidin conjugate or  $\text{R}\alpha\text{S}$  antiserum and HRPO-D $\alpha\text{R}$  antiserum. Colour changes were generated by incubation with ABTS and were monitored at 405 nm.

- 5 Only soluble pRAS112 protein (full length  $\text{ScFv}_{\text{NP}}$ -streptavidin monomer) can be detected in bacterial supernatants (Figure 12). Filtered bacterial growth medium recovered after induction of pRAS112 (⊙), pRAS113 (◆) or pRAS114 (■) protein was diluted in PBS and applied to wells of an ELISA plate coated with  $10 \mu\text{g}.\text{ml}^{-1}$  NIP<sub>15</sub>-BSA. Bound protein was
- 10 detected by serial incubation with Rabbit  $\alpha$  Streptavidin antisera and horseradish peroxidase conjugated Donkey  $\alpha$  Rabbit antisera, and colour changes generated by addition of ABTS were monitored at 405 nm. Like the parental  $\text{ScFv}_{\text{NP}}$ , this protein binds NIP<sub>15</sub>-BSA, but not lysozyme (Figure 13). Filtered bacterial growth medium recovered after induction
- 15 of pRAS112 protein was applied to wells of an ELISA plate coated with  $10 \mu\text{g}.\text{ml}^{-1}$  NIP<sub>15</sub>-BSA (⊙) or  $300 \mu\text{g}.\text{ml}^{-1}$  hen egg lysozyme (◆). Bound protein was detected by serial incubation with Rabbit  $\alpha$  Streptavidin antisera and horseradish peroxidase conjugated Donkey  $\alpha$  Rabbit antisera, and colour changes generated by addition of ABTS were monitored at 405
- 20 nm.

#### Partial purification of pRAS112 protein

- $\text{ScFv}_{\text{NP}}$ -streptavidin fusion protein (pRAS112 protein) can be concentrated
- 25 about 20-fold by precipitation from 50% saturated ammonium sulphate and dialysis against PBS. As expected concentrated pRAS112 protein binds iminobiotin-Sepharose at pH11, in contrast to parental  $\text{ScFv}_{\text{NP}}$  protein (Figure 14). Concentrated proteins resolubilised in PBS after precipitation from 50% (pRAS112) or 80% (pRAS111) saturated ammonium sulphate
  - 30 were applied at pH11 to a iminobiotin-Sepharose column (Pierce), and

antigen binding ability of material applied to the column (⊙) and material flowing through the column (♦) were measured by appropriate ELISA.

a) pRAS112 protein, b) pRAS111 protein.

5 The avidity of streptavidin fusions can be compared with univalent ScFvs.

a) The slope of a NIP-specific ELISA performed using pRAS112 streptavidin fusion differs from that performed using pRAS111 scFv.

10

b) pRAS112 protein binding to NIP-BSA cannot be competed with free NP, free NIP or NIP-BSA, whereas pRAS111 scFv can.

15

c) pRAS112 protein cannot be eluted in a single pulse from a NP-Sephacrose column. Multiple pulses of low pH interspersed with high pH washes are required to elute this protein. In contrast, pRAS111 scFv elutes with a single low pH step (Figure 34).

20

This indicates that the streptavidin fusions (pRAS112) are binding multivalently.

A representation of the pRAS112 protein is shown in Figure 15.

### Example 5: Construction of ScFv-BSRNase fusion molecules

25

#### Plasmid construction

30

Plasmids for the expression of ScFv-BSRNase fusions are shown in Figure 16. The plasmid pRAS111 is described in Example 1, and the plasmid pBSV5 is as described in Schein *et al*, *loc. cit*.



Figure 21 shows the sequence of the ScFv-BSRNase fusion (4-OH nitrophenacetyl antibody) inserted between the *Hind*III and *Eco*RI sites of plasmid pSP7 (available from Promega) to give plasmid pSPNPBSR as shown in Figure 16.

5

Figure 22 shows the sequence of the ScFv-BSRNase fusion (anti-human placental alkaline phosphatase antibody; H17E2) inserted between the *Hind*III and *Eco*RI sites of plasmid pSP71 to give plasmid pSPH17 $\Delta$ XBSR as shown in Figure 16.

10

The amino acid sequences of the V<sub>H</sub> and V<sub>L</sub> chains of H17E2 are disclosed in "Monoclonal Antibodies - applications in clinical oncology", pages 37-43, 1991, A.A. Epenetos, ed., Chapman & Hall, UK.

15

Figure 23 shows the sequence of the ScFv-BSRNase fusion (anti-lysozyme antibody) inserted between the *Hind*III and *Eco*RI sites of plasmid pUC18 (available from Pharmacia) to give pUCD1.3BSR as shown in Figure 16.

20

Figure 17 shows a diagrammatic representation of the specific case where a heterodimer has been synthesised and purified (as described *supra*), in this case each of the ScFvs recognises a different antigen on the same tumour cell.

25

The plasmids were made using standard methods of molecular biology as disclosed by Sambrook *et al* (1989) in *Molecular Cloning, a laboratory manual*, 2nd Edn, Cold Spring Harbor Laboratory Press, NY, USA.

30

The plasmid pSPNPBSR encodes a protein which directs cytotoxin RNase to a target cell-specific molecule derivatised with NP or NIP. The plasmid pPSH17 $\Delta$ XBSR encodes a protein which directs RNase to cells

expressing the human placental alkaline phosphatase antigen. The ScFv encoded by this plasmid is derived from the monoclonal antibody H17E2 (see Table 1).

- 5 In addition to the fusion gene consisting of the H17E2 scFv and seminal RNase only (see above) the following fusion genes which incorporate one or more of the following are useful:

- 10 (i) A C-terminal "KDEL" sequence (endoplasmic reticulum retention signal), which may elevate cytotoxicity by increasing the retention of the protein in the cell and reducing its loss to other endosomal pathways.
- (ii) A linker sequence at the N-terminus of the RNase to allow the N-terminus to be more flexible and increase the likelihood of forming
- 15 dimers.
- (iii) A disulphide loop containing sequence, derived from the diphtheria toxin, which allows the scFv and RNase to be linked via a disulphide bond, and permits efficient release of the RNase from the scFv once the cytotoxin has been internalised.

20

The plasmids which contain these genes (described diagrammatically in Figure 28 and individual nucleotide sequences encoding these proteins given in Figures 29 to 33) are identical to that expressing the original scFv-RNase fusion protein, ie only the DNA sequence of the actual

25 cytotoxic molecule has been altered. The conditions for expression and refolding are as described in the earlier Examples.

## Characterisation of the scFv-RNase protein

### *RNase activity of the fusion proteins.*

All the fusions described, H17-BSRNase, H17-DT-BSRNase, H17-DT-BSRNaseKDEL, H17-DT-Link-BSRNase, H17-DT-Link-BSRNaseKDEL, H17-BSRNaseKDEL, H17-Link-BSRNaseKDEL, have RNA-degrading activity, as demonstrated by an RNase assay which involves incubating a sample of the refolded protein (10-50 ng of crude fusion protein) with 5  $\mu$ g of RNA in a volume of 20  $\mu$ l at 37°C for 1 hr. In each case all the RNA was degraded, showing qualitative RNase activity in the fusion protein preparations.

### *Antigen-binding activity of fusion proteins.*

All the fusion proteins demonstrate binding to the antigen human placental alkaline phosphatase (hPLAP) in an ELISA system. The detecting layers for the ELISA were anti-bovine seminal RNase polyclonal antibodies (from rabbit) and anti-rabbit polyclonal antibodies (from goat).

### *Evidence for the dimeric nature of the scFv-RNase.*

Gel filtration experiments show the native molecular weight of the fusion proteins. Data from binding experiments indicates that the molecule has a higher avidity than the single-chain H17E2 antibody alone: the scFv will bind to an antigen affinity column (the antigen is placental alkaline phosphatase) and is eluted with a buffer consisting of 50 mM diethylamine (DEA), pH 12. The fusion protein, due to its higher avidity cannot be eluted under these mild conditions, and more harsh conditions are needed, eg 100 mM glycine, pH 2.2. Also, when the scFv and whole IgG H17E2 and fusion proteins are bound to their antigen on an ELISA plate and washed with copious amounts of 50 mM DEA, over 90% of the scFv is washed off, whereas only 40% of the whole IgG and fusion protein is

washed off. Finally, the shape of the ELISA curve for the whole IgG H17EE2 and fusion protein are similar (shallow slope), but that of the scFv is a steep slope. These experiments indicate that the scFv-RNase protein is dimeric.

5

*Cytotoxicity of the fusion proteins towards an antigen-positive cell-line (HEp2).*

HEp2 cells were seeded in 96-well microtitre plates and grown overnight in E4 culture media with 10 foetal calf serum (FCS) at a density of  $10^5$  cells per well. The next day, 10  $\mu$ l of crude refolded fusion protein in PBS was added to each well and allowed to grow for 72 hr. Cell-killing was detected using the Promega cell-titre 96 assay kit, which measures cell proliferation.

15 The scFv-BSRNase fusion protein consisting of a disulphide loop, KDEL and linker showed significant cell killing activity. The estimated final concentration of the cytotoxin was between 10-100 nM (see Figure 27 for the results of these experiments).

20 **Example 6: Construction of ScFv-DNaseI fusion molecules without a nuclear localization signal**

Plasmids for the expression of ScFv-DNaseI fusions are shown in Figure 18. The plasmid pRAS111 is described in Example 1 and  
25 M13mp19DNaseRec5 is described in Worrall and Connolly, *loc. cit.*

Figure 24 shows the sequence of the ScFv-DNaseI fusion (4-OH nitrophenacetyl antibody) inserted between the *Hind*III and *Bgl*III sites of plasmid pSP71 to give plasmid pSPNPDN1 as shown in Figure 18.

Figure 25 shows the sequence of the ScFv-DNaseI fusion (anti-human placental alkaline phosphatase antibody; H17E2) inserted between the *Hind*III and *Bgl*II sites of plasmid pSP71 to give plasmid pSPH17ΔXDN1 as shown in Figure 18.

5

Figure 26 shows the sequence of the ScFv-DNase fusion (anti-lysozyme antibody) inserted between the *Hind*III and *Bgl*II sites of plasmid pUC18 to give pUCD1.3DN1 as shown in Figure 18.

10 The plasmids were made using standard methods of molecular biology as disclosed in Sambrook *et al* (1989) in *Molecular Cloning, a laboratory manual*, 2nd Edn, Cold Spring Harbor Laboratory Press, NY, USA.

15 The plasmid pSPNPDN1 encodes a protein which directs DNaseI to a target cell-specific molecule derivatised with NP or NIP.

The plasmid pSPH17ΔXDN1 encodes a protein which directs DNaseI to cells expressing the human placental alkaline phosphatase antigen. The ScFv encoded by this plasmid is derived from the monoclonal antibody  
20 H17E2 (see Table 1).

The scFv-DNase I fusion has been expressed under identical conditions to that of the RNase fusions and refolded. The crude refolded preparation of the scFv-DNase I fusion protein shows PLAP-antigen binding activity  
25 in an ELISA system similar to the parent antibody H17E2. The detecting layers are anti-bovine DNase I (from rabbit) and anti-rabbit (from goat). The DNase I fusion protein also demonstrates DNA-degrading activity, in a similar system as that of the RNase assay, except 2 µg of DNA is incubated. The activity is only present when 10 mM CaCl<sub>2</sub> and 4 mM  
30 MgCl<sub>2</sub> is added, as is found with the naturally occurring bovine DNase I,

suggesting that functional scFv-DNase fusion molecules have been expressed and refolded from *E. coli*.

CLAIMS

1. A compound comprising a target cell-specific portion and a cytotoxic portion characterised in that the cytotoxic portion has nucleolytic activity.
2. A compound according to Claim 1 wherein the cytotoxic portion has ribonucleolytic activity.
3. A compound according to Claim 2 wherein the cytotoxic portion is a ribonuclease.
4. A compound according to Claim 3 wherein the ribonuclease is dimeric.
5. A compound according to Claim 4 wherein the ribonuclease is mammalian seminal ribonuclease.
6. A compound according to Claim 1 wherein the cytotoxic portion has DNA endonucleolytic activity.
7. A compound according to Claim 6 wherein the cytotoxic portion is at least the catalytically active portion of a DNA endonuclease.
8. A compound according to Claim 7 wherein the endonuclease is a mammalian deoxyribonuclease I.
9. A compound according to Claim 8 wherein a nuclear localization signal is incorporated.

10. A compound according to Claim 9 wherein the nuclear localization signal comprises the sequence PKKKRKV.
- 5 11. A compound according to Claim 7 wherein the DNA endonuclease is a restriction endonuclease.
12. A compound comprising a target cell-specific portion and a directly or indirectly cytotoxic second portion, characterised in that the target cell-specific portion recognises the target cell with high  
10 avidity.
13. A compound comprising a target cell-specific portion and a directly or indirectly cytotoxic portion characterised in that the cytotoxic portion is a sub-unit of an oligomer provided that, if the sub-unit is  
15 complexed with another sub-unit of the said oligomer then the said other sub-unit is the cytotoxic portion of a second compound of the invention.
14. A compound comprising an oligomeric complex of at least two  
20 molecules each comprising a target cell-specific portion and a further portion wherein the molecules are complexed to one another via their further portions.
15. A compound according to Claim 14 wherein the target cell-specific  
25 portion recognises a blood group antigen.
16. A compound comprising an oligomeric complex of at least two molecules each comprising a target cell-specific portion wherein the molecules are complexed to one another via their cytotoxic portions.



17. A compound according to Claim 16 wherein the respective cytotoxic portions of at least two of the molecules of the oligomeric complex are different from one another.
- 5 18. A compound according to Claim 16 wherein the respective target cell-specific portions of at least two of the molecules of the oligomeric complex are different from one another.
- 10 19. A compound comprising a target cell-specific portion and a directly or indirectly cytotoxic portion characterised in that the cytotoxic portion contains a binding site for a small molecule, wherein the said small-molecule binding site binds but does not modify catalytically the said small molecule.
- 15 20. A compound according to Claim 19 wherein the small-molecule binding site recognises the said small molecule with an affinity of at least  $K_D = 10^{-12}M$ .
- 20 21. A compound comprising a target cell-specific portion and a directly or indirectly cytotoxic portion characterised in that the target cell-specific portion comprises two or more binding sites for the target cell, wherein the target cell-specific portion is not an antibody or bivalent fragment thereof having respective arms which recognise the same entity as one another.
- 25 22. A compound according to any one of the preceding claims wherein the target cell-specific portion comprises an antibody or part thereof.
- 30 23. A compound according to any one of the preceding claims wherein the target cell-specific portion comprises a ScFv antibody fragment.

24. A compound according to any one of the preceding claims wherein the target cell-specific portion recognises and selectively binds to a tumour cell antigen.
- 5 25. A compound comprising a mediator portion and a directly or indirectly cytotoxic portion wherein the mediator portion recognises a target cell-specific molecule.
- 10 26. A compound according to Claim 25 wherein the target cell-specific molecule comprises a hapten and the mediator portion recognises the said hapten.
27. A compound according to Claim 26 wherein the said hapten is capable of chemical conjugation to the target cell-specific molecules.
- 15 28. A compound according to Claim 27 wherein the hapten is 4-hydroxy-3-nitrophenylacetic acid or 4-hydroxy-3-iodophenylacetic acid.
- 20 29. A compound according to any one of Claims 26 to 28 wherein the mediator is a ScFv.
30. A compound according to any one of Claims 26 to 29 wherein the cytotoxic portion comprises at least the biotin-binding portion of streptavidin.
- 25 31. A compound according to any one of Claims 26 to 29 wherein the cytotoxic portion has DNA endonucleolytic activity.
- 30 32. A compound according to Claim 31 wherein the cytotoxic portion

is at least the catalytically active portion of a DNA endonuclease.

- 5 33. A compound according to Claim 19 wherein the cytotoxic portion comprises at least the biotin- and subunit-binding portions of streptavidin.
- 10 34. A compound according to any of the preceding claims wherein the target cell-specific and cytotoxic portions of the compound are polypeptides.
35. A compound according to any of the preceding claims wherein the target cell-specific and cytotoxic portions of the compound are fused.
- 15 36. A nucleotide sequence encoding a compound according to Claim 35.
37. A therapeutic system comprising a compound according to Claim 19 and a drug, pro-drug, radionuclide, enzyme, antibody, toxin or any other reagent which is conjugated to the said small molecule.
- 20 38. A therapeutic system according to Claim 37 wherein the said small molecule is biotin.
39. A therapeutic system comprising a compound according to Claim 25 and a target cell-specific molecule.
- 25 40. A therapeutic system comprising a compound according to Claim 30, a target cell-specific molecule and a drug, pro-drug, radionuclide, enzyme, antibody toxin or any other reagent which is conjugated to biotin.
- 30

41. A pharmaceutical composition comprising a compound according to any one of Claims 1 to 35 and a pharmaceutical carrier.
- 5 42. A method of treating a mammal having target cells to be destroyed, the method comprising (1) administering a compound according to any one of Claims 12, 13 and 16 to 21, wherein the cytotoxic portion is indirectly cytotoxic to the mammal, (2) allowing the ratio of (compound bound to the target cells) : (compound not bound to the target cells) to reach a desired value, and (3) administering a  
10 drug, pro-drug, radionuclide, enzyme, antibody, toxin or any other reagent which is conjugated to a small molecule.
43. A method of treating a mammal according to Claim 42 wherein the said small molecule is biotin.  
15
44. A method of treating a mammal having target cells to be destroyed, the method comprising administering the compound according to any one of Claims 1 to 13 and 16 to 24 wherein the cytotoxic portion is directly cytotoxic.  
20
45. A method of treating a mammal having target cells to be destroyed, the method comprising administering a target cell-specific molecule. (2) allowing the ratio of (molecule bound to the target cells):(molecule not bound to the target cells) to reach a desired  
25 value, (3) administering a compound according to Claim 30 wherein the cytotoxic portion is indirectly cytotoxic, (4) allowing the ratio of (compound bound to target cell-specific molecule):(compound not bound to target cell-specific molecule) to reach a desired value and (5) administering a drug, pro-drug, radionuclide, enzyme, antibody,  
30 toxin or any other reagent which is conjugated to biotin.

46. A method of treating a mammal having target cells to be destroyed, the method comprising administering a target cell-specific molecule. (2) allowing the ratio of (molecule bound to target cells):(molecule not bound to target cells) to reach a desired value, and (3) administering a compound according to any one of Claims 25 to 29 wherein the cytotoxic portion is directly cytotoxic.
- 5
47. A method of determining a blood group of an individual, the method comprising (1) removing blood from an individual into a container, (2) adding a compound according to Claim 15 and (3) observing the extent of haemagglutination.
- 10

**Amendments to the claims have been filed as follows**

1. A compound comprising a target cell-specific portion fused to an oligomeric ribonuclease.  
5
2. A compound according to Claim 1 wherein the ribonuclease is mammalian seminal ribonuclease.
3. A compound according to Claim 1 wherein the target cell-specific  
10 portion comprises a ScFv.
4. A compound according to any one of the preceding claims wherein each subunit of the said ribonuclease has a target cell-specific portion fused thereto.  
15
5. A compound according to any one of the preceding claims wherein the target cell-specific portion bind selectively to a tumour cell.
6. A nucleic acid encoding a compound as defined in any one of  
20 Claims 1 to 4.
7. A compound comprising an oligomeric complex of at least two molecules each comprising a target cell-specific portion fused to a further portion wherein the molecules are complexed to one  
25 another via their further portions and wherein the target cell-specific portion and the further portion are not derived from the same molecule.
8. A compound comprising a target cell-specific portion and a further  
30 portion characterised in that the target cell-specific portion

comprises two or more binding sites for the target cell, wherein the target cell-specific portion is not an antibody or bivalent fragment thereof having respective arms which recognise the same entity as one another.

5

9. A compound according to Claim 7 or 8 wherein the further portion is directly or indirectly cytotoxic.

10

10. A compound according to any one of Claims 7 or 9 wherein the further portion is a subunit of an oligomeric polypeptide.

11. A compound according to Claim 10 wherein the oligomeric polypeptide is mammalian seminal ribonuclease or at least the biotin- and subunit-binding portions of streptavidin.

15

12. A compound according to Claim 7 or 8 wherein the target cell-specific portion recognises a blood group antigen.

20

13. A compound according to any one of Claims 7 or 12 wherein the target cell-specific portion comprises a ScFv antibody fragment.

14. A compound according to Claim 8 wherein the target cell-specific portion and the further portion are fused.

25

15. A nucleotide sequence encoding a compound according to Claim 7 or 14.

30

16. A compound comprising a portion which recognises 4-hydroxy-3-nitrophenylacetic acid or 4-hydroxy-3-iodophenylacetic acid and a directly or indirectly cytotoxic portion.

17. A compound according to Claim 16 wherein the recognition portion is a ScFv.
18. A compound according to Claim 16 or 17 wherein the cytotoxic portion comprises at least the biotin-binding portion of streptavidin.
19. A compound according to Claim 16 or 17 wherein the cytotoxic portion has nucleolytic activity.
20. A compound according to Claim 19 wherein the cytotoxic portion is at least the catalytically active portion of a DNA endonuclease.
21. A compound according to Claim 19 wherein the cytotoxic portion is at least the catalytically active portion of a RNase.
22. A therapeutic system comprising a compound according to Claim 16 and a target cell-specific molecule with 4-hydroxy-3-nitrophenylacetic acid or 4-hydroxy-3-iodophenylacetic acid conjugated thereto.
23. A therapeutic system comprising a compound according to Claim 18, a target cell-specific molecule and a drug, pro-drug, radionuclide, enzyme, antibody toxin or any other reagent which is conjugated to biotin.
24. A pharmaceutical composition comprising a compound according to any one of Claims 1 to 21 and a pharmaceutical carrier.
25. A method of treating a mammal having target cells to be destroyed, the method comprising administering the compound according to any



one of Claims 1 to 5 and 7 to 14 wherein the cytotoxic portion is directly cytotoxic.

26. A method of treating a mammal having target cells to be destroyed,  
5 the method comprising (1) administering a compound according to  
any one of Claims 7 or 14, wherein the cytotoxic portion is  
indirectly cytotoxic to the mammal, (2) allowing the ratio of  
(compound bound to the target cells) : (compound not bound to the  
10 target cells) to reach a desired value, and (3) administering, as  
appropriate, a drug, pro-drug, radionuclide, enzyme, antibody,  
toxin, or any other reagent which is conjugated to a small molecule.
27. A method of treating a mammal according to Claim 26 wherein the  
15 said small molecule is biotin.
28. A method of determining a blood group of an individual, the method  
comprising (1) removing blood from an individual into a container,  
(2) adding a compound according to Claim 12 and (3) observing the  
20 extent of haemagglutination.



The  
Patent  
Office  
74

Application No: GB 9615814.2  
Claims searched: 1 to 24, 28

Examiner: Colin Sherrington  
Date of search: 10 September 1996

**Patents Act 1977**  
**Search Report under Section 17**

**Databases searched:**

UK Patent Office collections, including GB, EP, WO & US patent specifications, in:

UK Cl (Ed.O): C3H(HB7E,HB7M,HB7P)

Int Cl (Ed.6): A61K 47/48

Other: ONLINE: WPI

**Documents considered to be relevant:**

Category	Identity of document and relevant passage	Relevant to claims
X	WO 91/16069 A1 (THE UNITED STATES OF AMERICA, as represented by THE SECRETARY, U.S. DEPARTMENT OF COMMERCE) -whole document	1 to 7, 9 to 14, 24, 28
A	Cell Biophysics 1992,21,121-138 -Susanna M.Rybak et al. "Rational Immunotherapy with Ribonuclease Chimeras: An Approach Toward Humanizing Immunotoxins"	1 to 5

X Document indicating lack of novelty or inventive step  
Y Document indicating lack of inventive step if combined with one or more other documents of same category.  
& Member of the same patent family

A Document indicating technological background and/or state of the art.  
P Document published on or after the declared priority date but before the filing date of this invention.  
E Patent document published on or after, but with priority date earlier than, the filing date of this application.



# (12) UK Patent Application (19) GB (11) 2 300 859 (13) A

(43) Date of A Publication 20.11.1996

(21) Application No 9615814.2

(22) Date of Filing 17.01.1994

Date Lodged 26.07.1996

(30) Priority Data

(31) 9300686 (32) 15.01.1993 (33) GB

(62) Derived from Application No. 9513418.5  
under Section 15(4) of the Patents Act 1977

(71) Applicant(s)

Imperial Cancer Research Technology Limited

(Incorporated in the United Kingdom)

Sardinia House, Sardinia Street, LONDON,  
WC2A 3NL, United Kingdom

(72) Inventor(s)

Agamemnon Antoniou Epenetos  
Robert Anthony Spooner  
Mahendra Deonarain

(51) INT CL<sup>6</sup>

A61K 47/48, C07K 16/00 19/00, C12N 15/62, G01N  
33/80, // A61K 51/10, C12N 9/22 (C07K 19/00 C07M  
5:00)

(52) UK CL (Edition O)

C3H HB7M HB7P H650 H651 H654 H656 H657 H660  
H675 H684 H685 H686 H690 H730  
U1S S1313 S1332 S1333 S1334 S1337

(56) Documents Cited

WO 91/16069 A1  
Cell Biophysics 1992,21,121-138

(58) Field of Search

UK CL (Edition O) C3H HB7E HB7M HB7P  
INT CL<sup>6</sup> A61K 47/48  
ONLINE: WPI

(74) Agent and/or Address for Service

Eric Potter Clarkson  
St Mary's Court, St Mary's Gate, NOTTINGHAM,  
NG1 1LE, United Kingdom


(54) Compounds comprising a target cell-specific portion (TCP) and a cytotoxic portion (CP)

(57) Compounds are described which are selected from:

- (i) TCP, preferably an ScFv and/or one which binds selectively to a tumour cell, fused to an oligomeric ribonuclease, preferably mammalian seminal ribonuclease;
- (ii) a oligomeric complex of at least two molecules, each comprising a TCP fused to a further (cytotoxic) portion, said molecules being complexed to one another via said further portion;
- (iii) TCP, other than an antibody, comprising at least two binding sites for the target cell, linked to a further (cytotoxic) portion;
- (iv) a recognition portion, preferably an ScFv, for 4-hydroxy-3-iodophenylacetic acid or 4-hydroxy-3-nitrophenylacetic acid bonded to CP, preferably the biotin-binding portion of streptavidin or one which has nucleolytic activity, especially the catalytic activity of either DNA endonuclease or RNAase, which may be bonded to TCP.

In compounds (ii) and (iii), the further (cytotoxic) portion may be mammalian seminal ribonuclease or a biotin-binding portion of streptavidin and the TCP preferably recognises either a blood group antigen (for use in the determination of a blood group by haemagglutination) or an ScFv antibody fragment. Nucleic acid sequences encoding compounds (i) to (iii) are disclosed.

The compounds are of use in therapeutic systems for the treatment of cancer.



GB 2 300 859 A